

REPRESENTATION TYPE OF FINITE QUIVER HECKE ALGEBRAS OF TYPE $A_{2\ell}^{(2)}$

SUSUMU ARIKI¹ AND EUIYONG PARK²

ABSTRACT. We study cyclotomic quiver Hecke algebras $R^{\Lambda_0}(\beta)$ in type $A_{2\ell}^{(2)}$, where Λ_0 is the fundamental weight. The algebras are natural $A_{2\ell}^{(2)}$ -type analogue of Iwahori-Hecke algebras associated with the symmetric group, from the viewpoint of the Fock space theory developed by the first author and his collaborators. We give a formula for the dimension of the algebra, and a simple criterion to tell the representation type. The criterion is a natural generalization of Erdmann and Nakano's for the Iwahori-Hecke algebras. Except for the examples coming from cyclotomic Hecke algebras, no results of these kind existed for cyclotomic quiver Hecke algebras, and our results are the first instances beyond the case of cyclotomic Hecke algebras.

INTRODUCTION

Block algebras of the finite Hecke algebra associated with the symmetric group have been studied extensively in 1990's. Let \mathbf{k} be an algebraically closed field, $q \in \mathbf{k}^\times$ the parameter of the Hecke algebra, and e the *quantum characteristic* defined by

$$e = \min\{a \in \mathbb{Z}_+ \mid 1 + q + \cdots + q^{a-1} = 0 \text{ holds in } \mathbf{k}\}.$$

Then, the block algebras are parametrized by pairs of an e -core partition and an e -weight. This well-known fact, the *Nakayama conjecture* for Hecke algebras, was proved by James and Mathas [12, Thm. 4.29] based on earlier work by Dipper and James [7]. Nowadays it is understood as one of the results which fits in Fock space theory for cyclotomic Hecke algebras: the block algebras categorify weight spaces of the basic $\mathfrak{g}(A_{e-1}^{(1)})$ -module $V(\Lambda_0)$. In fact, this example was the origin of the conjecture by Lascoux, Leclerc and Thibon, where the Fock space theory started. Let δ be the null root and let W be the Weyl group i.e. the affine symmetric group. We recall that every weight that appear in $V(\Lambda_0)$ has the form $w\Lambda_0 - k\delta$, where $w \in W$ and $k \in \mathbb{Z}_{\geq 0}$. Then, in the Misra-Miwa realization of the Fock space, $w\Lambda_0$ is the e -core partition, and k is the e -weight. Thus, the modular representation theory of finite Hecke algebras turned out to be Lie theoretic, and we have various interesting questions to

¹ This work is supported in part by JSPS, Grant-in-Aid for Scientific Research (B) 23340006.

² This work is supported by JSPS Postdoctoral Fellowship for Foreign Researchers.

ask under the philosophy of categorification. For example, research in the early stage focused on higher level analogues, that is, the modular representation theory of cyclotomic Hecke algebras, and finite Hecke algebras of type BC (and type D by Clifford theory) are special cases of them. See [1], for example.

An important turning point for the philosophy was the introduction of cyclotomic quiver Hecke algebras by Khovanov and Lauda in [26], [27]. The algebras are associated with a pair of dominant integral weight Λ and non-negative integral linear combination β of simple roots. We denote the algebra by $R^\Lambda(\beta)$. As Rouquier [41] also developed another theory to categorify integrable highest weight modules, which generalizes his work with Chuang [5], it is also called the *cyclotomic KLR algebra* by various authors. Further, theorems from Brundan and Kleshchev [4] and Lyle and Mathas [35] combined imply that block algebras of the cyclotomic Hecke algebras are cyclotomic quiver Hecke algebras of type $A_{e-1}^{(1)}$. More precisely, we label the vertices of the Dynkin diagram with $\mathbb{Z}/e\mathbb{Z}$ as usual, and we let $A = (a_{ij})_{i,j \in \mathbb{Z}/e\mathbb{Z}}$ be the Cartan matrix. Then their quiver Hecke algebra is associated with polynomials

$$Q_{i,j}(u, v) = 1, \text{ if } a_{ij} = 0, \text{ and } Q_{i,i+1}(u, v) = Q_{i+1,i}(v, u) = -(u - v)^{-a_{i,i+1}}, \text{ for } i \in \mathbb{Z}/e\mathbb{Z}.$$

Therefore, the cyclotomic quiver Hecke algebras are indeed generalization of cyclotomic Hecke algebras.

Many papers on the quiver Hecke algebras have already appeared, e.g. [2], [10], [21], [28], [29], [33] or [42]. But most of them study labeling or construction of irreducible modules. The only case which receives rather detailed study is type $A_{e-1}^{(1)}$, where the Brundan-Kleshchev isomorphism theorem allows us to reduce problems to well-studied cyclotomic Hecke algebras.

Our aim in this paper is to show some new results which go beyond this stage, and we study $R^{\Lambda_0}(\beta)$ in type $A_{2\ell}^{(2)}$ in our paper. As we have explained in the above, the algebra is a natural $A_{2\ell}^{(2)}$ -type analogue of the Iwahori-Hecke algebra associated with the symmetric group. Thus, we call them *finite quiver Hecke algebras of type $A_{2\ell}^{(2)}$* . We give dimension formula for $R^{\Lambda_0}(\beta)$, some structure theorem, and prove Erdmann-Nakano type theorem which tells the representation type of $R^{\Lambda_0}(\beta)$. Note that the Erdmann-Nakano theorem treats $e = 2$ case separately, and tame representation type appears only in this case. The reader might expect that the statement in type $A_{2\ell}^{(2)}$ would be different from the other $A_{2\ell}^{(2)}$, for $\ell \geq 2$. However, we find that it is not the case, and the criterion is uniform for all ℓ . Apart this point, the criterion is exactly the same as that of the Erdmann-Nakano theorem, for $e \geq 3$, as follows.

Theorem A (Theorem 4.22). Suppose that $R^{\Lambda_0}(\beta)$ is a finite quiver Hecke algebra of type $A_{2\ell}^{(2)}$, for $\ell \geq 1$. We denote the Weyl group of type $A_{2\ell}^{(2)}$ by W , $W\Lambda_0$ the W -orbit through Λ_0 . There exist unique $\kappa \in W\Lambda_0$ and unique $k \in \mathbb{Z}_{\geq 0}$ such that $\beta = \Lambda_0 - \kappa + k\delta$. Then, $R^{\Lambda_0}(\beta)$ is

- (1) simple if $k = 0$,
- (2) of finite representation type but not semisimple if $k = 1$,
- (3) of wild representation type if $k \geq 2$,
- (4) and tame representation type does not occur.

The following is our dimension formula. We do not explain the notation here. Note that almost no result is known for the dimension of cyclotomic quiver Hecke algebras, except for block algebras of cyclotomic Hecke algebras. Another importance of Theorem B is that it tells when $e(\nu) = 0$, for $\nu \in I^n$. It is repeatedly used in the proof of Theorem A.

Theorem B (Theorem 3.4). Let λ be a shifted partition, $\nu \in I^n$. Then, for $\beta \in \mathbb{Q}^+$ and $\nu, \nu' \in I^\beta$, we have

$$\begin{aligned} \dim e(\nu') R^{\Lambda_0}(n) e(\nu) &= \sum_{\lambda \vdash n} 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} K(\lambda, \nu') K(\lambda, \nu), \\ \dim R^{\Lambda_0}(\beta) &= \sum_{\lambda \vdash n, \text{wt}(\lambda) = \Lambda_0 - \beta} 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} |\text{ST}(\lambda)|^2, \\ \dim R^{\Lambda_0}(n) &= \sum_{\lambda \vdash n} 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} |\text{ST}(\lambda)|^2. \end{aligned}$$

In proving our results, solution to the cyclotomic categorification conjecture from [26] by Kang and Kashiwara in [17] and [24] is used in an essential way. The strategy of the proof works for general affine types. In particular, the argument in our paper gives a completely new proof of the original Erdmann-Nakano theorem [8, 1.2] as well. We will consider the other affine types than $A_{e-1}^{(1)}$ and $A_{2\ell}^{(2)}$ in a separate paper.

The structure of the paper is as follows. Section 1 is for preliminaries. In section 2, we recall cyclotomic quiver Hecke algebras and categorification results. In section 3, we prove the dimension formula. In section 4, we prove the Erdmann-Nakano type theorem for the representation type of $R^{\Lambda_0}(\beta)$. Section 5 is an appendix for generalized cellularity. We propose a generalized cellular algebra structure for finite quiver Hecke algebras of type $A_{2\ell}^{(2)}$. We are grateful to Mr. Mori for letting us know his work on generalized cellularity and some comments on our paper.

1. PRELIMINARIES

1.1. Shifted Young diagrams. An array λ of a finite number of boxes arranged in l rows is a *shifted Young diagram of depth l* , if

- (i) the i th row from the top, for $1 \leq i \leq l$, starts with its leftmost box in the i th column and there is no gap before it ends on the rightmost box,
- (ii) each row has strictly shorter length than its predecessor.

We denote the depth l by $l(\lambda)$ and we denote $\lambda \vdash n$ if λ consists of n boxes. We write $(i, j) \in \lambda$ if there exists a box in the i th row and the j th column. We identify a shifted Young diagram λ with the strict partition $(\lambda_1 > \lambda_2 > \lambda_3 > \cdots)$, where λ_i is the number of boxes in the i th row of λ . For $\lambda = (\lambda_1 > \lambda_2 > \cdots) \vdash n$ and $\mu = (\mu_1 > \mu_2 > \cdots) \vdash n$, we say λ *dominates* μ and write $\lambda \succeq \mu$ if

$$(1.1.1) \quad \sum_{i=1}^k \lambda_i \geq \sum_{i=1}^k \mu_i, \text{ for } 1 \leq k \leq l(\lambda).$$

A *standard tableau* T of shape $\lambda \vdash n$ is a filling of n boxes of λ with numbers 1 to n such that (i) each number is used exactly once, (ii) the numbers in rows and columns increase from left to right and top to bottom, respectively. Let $\text{ST}(\lambda)$ be the set of all standard tableaux of shape λ . For example, there are 3 standard tableaux of shape $(4, 1)$:

$$\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline & 5 & & \\ \hline \end{array}, \quad \begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 5 \\ \hline & 4 & & \\ \hline \end{array}, \quad \begin{array}{|c|c|c|c|} \hline 1 & 2 & 4 & 5 \\ \hline & 3 & & \\ \hline \end{array}.$$

For $\lambda = (\lambda_1 > \lambda_2 > \cdots) \vdash n$, let T^λ be the tableau of shape λ whose (i, j) -entry is

$$(j - i) + 1 + \sum_{k=1}^{i-1} \lambda_k.$$

We call T^λ the *canonical tableau* of shape λ . In the above example, the first tableau is the canonical tableau of shape $(4, 1)$. The following theorem is well-known.

Theorem 1.1. (1) *For $\lambda \vdash n$, the number of standard tableaux of shape λ is given by the following hook formula.*

$$|\text{ST}(\lambda)| = \frac{n!}{\prod_{(i,j) \in \lambda} h_{i,j}},$$

where $h_{i,j}$ is the hook-length of $(i, j) \in \lambda$, i.e.,

$$h_{i,j} = |\{(i, j') \in \lambda \mid j' \geq j\}| + |\{(i', j) \in \lambda \mid i' > i\}| + |\{(j+1, j') \in \lambda \mid j' > j\}|.$$

(2) *We have the following equality.*

$$n! = \sum_{\lambda \vdash n} 2^{n-l(\lambda)} |\text{ST}(\lambda)|^2.$$

Let $\lambda \vdash n$ be a shifted Young diagram. We declare that each row has the residue pattern

$$0 \ 1 \ 2 \ \cdots \ \ell \ \cdots \ 2 \ 1 \ 0,$$

which repeats from left to right in each row, and define $\text{res}(i, j)$ to be the residue of $(i, j) \in \lambda$. For example, if \mathfrak{g} is of type $A_6^{(2)}$ and $\lambda = (10, 6, 3, 1)$, the residues are as follows:

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| 0 | 1 | 2 | 3 | 2 | 1 | 0 | 0 | 1 | 2 |
| | 0 | 1 | 2 | 3 | 2 | 1 | | | |
| | | 0 | 1 | 2 | | | | | |
| | | | 0 | | | | | | |

Thus, if we pick $(2, 5) \in \lambda$ then it has residue 3. Now, we are ready to introduce the residue sequence of a tableau.

Definition 1.2. Let $T \in \text{ST}(\lambda)$. We define the *residue sequence* of T by

$$\text{res}(T) = (\text{res}(i_1, j_1), \text{res}(i_2, j_2), \dots, \text{res}(i_n, j_n)) \in I^n,$$

where $(i_k, j_k) \in \lambda$ is the box filled with k in T , for $1 \leq k \leq n$.

We remark that the residue pattern for shifted Young diagrams is the same as the pattern of Young walls defined by the level one perfect crystal of type $A_{2\ell}^{(2)}$ [11, 16, 18, 19], which we will use in Section 2.3. It reminds the reader of the q -deformed Fock space introduced in [20], and of the q -deformed Fock space from [25]. However, we must distinguish between categorification of integrable highest weight modules and its crystals. It seems more natural to consider the embedding of the basic module $V(\Lambda_0)$ into the classical Fock space which we will recall in Section 1.3, rather than the q -deformed Fock space. We suspect that some theory of packets would explain the distinction.

1.2. Cartan datum. Let $I = \{0, 1, \dots, \ell\}$, for $\ell \geq 2$, and let A be the *affine Cartan matrix* of type $A_{2\ell}^{(2)}$, i.e.,

$$A = (a_{ij})_{i,j \in I} = \begin{pmatrix} 2 & -2 & 0 & \dots & 0 & 0 & 0 \\ -1 & 2 & -1 & \dots & 0 & 0 & 0 \\ 0 & -1 & 2 & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 2 & -1 & 0 \\ 0 & 0 & 0 & \dots & -1 & 2 & -2 \\ 0 & 0 & 0 & \dots & 0 & -1 & 2 \end{pmatrix}.$$

When $\ell = 1$, the affine Cartan matrix of type $A_2^{(2)}$ is

$$A = (a_{ij})_{i,j \in I} = \begin{pmatrix} 2 & -4 \\ -1 & 2 \end{pmatrix}.$$

We choose a realization of the Cartan matrix and obtain an *affine Cartan datum* (A, P, Π, Π^\vee) , where

- (1) A is the affine Cartan matrix of type $A_{2\ell}^{(2)}$ as above,
- (2) P is a free abelian group of rank $\ell + 2$, called the *weight lattice*,
- (3) $\Pi = \{\alpha_i \mid i \in I\} \subset P$, called the set of *simple roots*,
- (4) $\Pi^\vee = \{h_i \mid i \in I\} \subset P^\vee := \text{Hom}(P, \mathbb{Z})$, called the set of *simple coroots*,

which satisfy the following properties:

- (a) $\langle h_i, \alpha_j \rangle = a_{ij}$ for all $i, j \in I$,
- (b) Π and Π^\vee are linearly independent, respectively.

The weight lattice P has a symmetric bilinear pairing (\mid) satisfying

$$(\alpha_i \mid \Lambda) = d_i \langle h_i, \Lambda \rangle \quad \text{for all } \Lambda \in P,$$

where $(d_0, d_1, \dots, d_\ell) = (1, 2, \dots, 2, 4)$. We denote by

$$P^+ = \{\Lambda \in P \mid \Lambda(h_i) \in \mathbb{Z}_{\geq 0}, i \in I\}$$

the set of *dominant integral weights*. We fix an element d which satisfies $\langle d, \alpha_i \rangle = \delta_{i0}$. It is called the *scaling element*. Then, for $i \in I$, we define the i th *fundamental weight* $\Lambda_i \in P^+$, i.e. the element defined by $\Lambda_i(d) = 0$ and $\Lambda_i(h_j) = \delta_{ij}$, for $j \in I$. The free abelian group $Q = \bigoplus_{i \in I} \mathbb{Z}\alpha_i$ is called the *root lattice*, and $Q^+ = \sum_{i \in I} \mathbb{Z}_{\geq 0}\alpha_i$ is the *positive cone* of the root lattice. For $\beta = \sum_{i \in I} k_i \alpha_i \in Q^+$, the *height* of β is defined by $|\beta| = \sum_{i \in I} k_i$.

Let $W = \langle r_i \mid i \in I \rangle$ be the *Weyl group* associated with A . The Coxeter generators $\{r_i\}_{i \in I}$ act on P by $r_i \Lambda = \Lambda - \langle h_i, \Lambda \rangle \alpha_i$, for $\Lambda \in P$, as usual. Finally, the *null root* in type $A_{2\ell}^{(2)}$ is

$$\delta = 2\alpha_0 + 2\alpha_1 + \dots + 2\alpha_{\ell-1} + \alpha_\ell,$$

for $\ell \geq 1$. Note that $\langle h_i, \delta \rangle = 0$ and $w\delta = \delta$, for $i \in I$ and $w \in W$. The residue pattern for shifted Young diagrams comes from the null root, and we denote the residue pattern $01 \dots \ell \dots 10$ by ν_δ .

1.3. The Fock space of neutral fermions. Recall that $\mathfrak{g}(A)$ is realized as $BKP_{2\ell+1}$ in the terminology of [6, §2]. In other words, $\mathfrak{g}(A)$ is obtained by $(2\ell+1)$ -reduction from $\mathfrak{g}(B_\infty)$. To obtain the basic $\mathfrak{g}(A)$ -module $V(\Lambda_0)$, we start with the basic $\mathfrak{g}(B_\infty)$ -module. Thus, we recall the Fock space of neutral fermions from [14, §6], which affords the basic spin representation of $\mathfrak{g}(B_\infty)$. Let C be the Clifford algebra over complex numbers \mathbb{C} defined by generators ϕ_k ($k \in \mathbb{Z}$) and the following relations:

$$\phi_p \phi_q + \phi_q \phi_p = \begin{cases} 2 & \text{if } p = q = 0, \\ (-1)^p \delta_{p, -q} & \text{otherwise.} \end{cases}$$

The Chevalley generators of $\mathfrak{g}(B_\infty)$ are given as follows:

$$\begin{aligned} \mathbf{e}_0 &= \phi_{-1}\phi_0, & \mathbf{e}_j &= (-1)^j \phi_{-j-1}\phi_j & (j \geq 1), \\ \mathbf{f}_0 &= \phi_1\phi_0, & \mathbf{f}_j &= (-1)^j \phi_{j+1}\phi_{-j} & (j \geq 1), \\ \mathbf{h}_0 &= 2\phi_1\phi_{-1} + 1, & \mathbf{h}_j &= (-1)^j (\phi_j\phi_{-j} + \phi_{j+1}\phi_{-j-1}) & (j \geq 1). \end{aligned}$$

Let \mathbf{l} be the left ideal of \mathbb{C} generated by $\{\phi_k \mid k < 0\}$, and define $\mathbf{F} = \mathbb{C}/\mathbf{l}$. Then, \mathbf{F} is the direct sum of two irreducible $\mathfrak{g}(B_\infty)$ -modules $\mathbf{F}(\Lambda_0)$ and $\mathbf{F}(\Lambda_1)$, and their highest weight vectors are $|0\rangle := 1 + \mathbf{l}$ and $|1\rangle := \phi_0|0\rangle$, respectively. We take a shifted Young diagram λ . It can be written as $\lambda = (\lambda_1 > \lambda_2 > \dots > \lambda_{2r-1} > \lambda_{2r} \geq 0)$ for a unique r , where $\lambda_{2r} > 0$ if $\ell(\lambda)$ is even and $\lambda_{2r} = 0$ if $\ell(\lambda)$ is odd. We define $|\lambda\rangle = \phi_{\lambda_1}\phi_{\lambda_2} \cdots \phi_{\lambda_{2r}}|0\rangle$. Then, they are linearly independent and

$$\mathcal{F} := \mathbf{F}(\Lambda_0) = \text{Span}_{\mathbb{C}}\{|\lambda\rangle \mid \lambda : \text{shifted Young diagrams}\}.$$

It is easy to see that if λ has a row of length $j+1$ and does not have a row of length j (resp. λ has a row of length j and does not have a row of length $j+1$), then we have

$$(1.3.1) \quad \mathbf{e}_j|\lambda\rangle = |\mathbf{e}_j\lambda\rangle, \quad (\text{resp. } \mathbf{f}_j|\lambda\rangle = |\mathbf{f}_j\lambda\rangle),$$

where $\mathbf{e}_j\lambda$ (resp. $\mathbf{f}_j\lambda$) is the shifted Young diagram obtained from λ by deleting the rightmost box of the row of length $j+1$ (resp. adding a new box on the right of the row of length j). Otherwise, we have $\mathbf{e}_j|\lambda\rangle = 0$ (resp. $\mathbf{f}_j|\lambda\rangle = 0$). Therefore, we may think of \mathcal{F} as a based vector space whose basis is given by $|\lambda\rangle$'s, and it is an integrable $\mathfrak{g}(B_\infty)$ -module by the action $|\lambda\rangle \mapsto |\mathbf{e}_j\lambda\rangle, |\mathbf{f}_j\lambda\rangle$. This is our *Fock space*.

We now define a \mathfrak{g} -module structure on the space \mathcal{F} by reduction. Let $h = 2\ell + 1$. From [6, Table 2], the action of the Chevalley generators f_i, e_i of \mathfrak{g} on \mathcal{F} are given as follows:

$$(1.3.2) \quad f_i = \sum_{j \geq 0, j \equiv i, -i-1} \mathbf{f}_j, \quad e_i = \begin{cases} \mathbf{e}_0 + 2 \sum_{j > 0, j \equiv 0, -1} \mathbf{e}_j & \text{if } i = 0, \\ \sum_{j > 0, j \equiv i, -i-1} \mathbf{e}_j & \text{if } i = 1, \dots, \ell, \end{cases}$$

where all congruences are taken modulo h . Each $|\lambda\rangle$ is a weight vector, and if we define a multiset $\text{res}(\lambda)$ of I by $\text{res}(\lambda) = \{\text{res}(i, j) \mid (i, j) \in \lambda\}$, then its weight is given by

$$\text{wt}(\lambda) = \Lambda_0 - \sum_{k \in \text{res}(\lambda)} \alpha_k.$$

We remark that \mathcal{F} is isomorphic to the direct sum of countably many copies of $V(\Lambda_0)$.

2. CYCLOTOMIC QUIVER HECKE ALGEBRAS OF TYPE $A_{2\ell}^{(2)}$

2.1. Quiver Hecke algebras. Throughout the paper, \mathbf{k} is an algebraically closed field, and algebras are unital associative \mathbf{k} -algebras.

Let (A, P, Π, Π^\vee) be the affine Cartan datum from 1.2. We are going to define \mathbf{k} -algebras which we call *finite quiver Hecke algebras of type $A_{2\ell}^{(2)}$* . To do this, we choose polynomials $\mathcal{Q}_{i,j}(u, v) \in \mathbf{k}[u, v]$, for $i, j \in I$, of the form

$$\mathcal{Q}_{i,j}(u, v) = \begin{cases} \sum_{p(\alpha_i|\alpha_i)+q(\alpha_j|\alpha_j)+2(\alpha_i|\alpha_j)=0} t_{i,j;p,q} u^p v^q & \text{if } i \neq j, \\ 0 & \text{if } i = j, \end{cases}$$

where $t_{i,j;p,q} \in \mathbf{k}$ are such that $t_{i,j;-a_{ij},0} \neq 0$ and $\mathcal{Q}_{i,j}(u, v) = \mathcal{Q}_{j,i}(v, u)$.

The symmetric group $\mathfrak{S}_n = \langle s_k \mid k = 1, \dots, n-1 \rangle$ acts on I^n by place permutations. Namely, the Coxeter generator s_k acts on I^n by

$$s_k(\nu_1, \dots, \nu_k, \nu_{k+1}, \dots, \nu_n) = (\nu_1, \dots, \nu_{k+1}, \nu_k, \dots, \nu_n).$$

Definition 2.1. Let $\Lambda \in P^+$. The *cyclotomic quiver Hecke algebra* $R^\Lambda(n)$ associated with polynomials $(\mathcal{Q}_{i,j}(u, v))_{i,j \in I}$ and the dominant integral weight Λ is the \mathbb{Z} -graded \mathbf{k} -algebra defined by three sets of generators

$$\{e(\nu) \mid \nu = (\nu_1, \dots, \nu_n) \in I^n\}, \{x_k \mid 1 \leq k \leq n\}, \{\psi_l \mid 1 \leq l \leq n-1\}$$

subject to the following relations:

$$\begin{aligned} e(\nu)e(\nu') &= \delta_{\nu,\nu'} e(\nu), \quad \sum_{\nu \in I^n} e(\nu) = 1, \quad x_k e(\nu) = e(\nu) x_k, \quad x_k x_l = x_l x_k, \\ \psi_l e(\nu) &= e(s_l(\nu)) \psi_l, \quad \psi_k \psi_l = \psi_l \psi_k \text{ if } |k-l| > 1, \\ \psi_k^2 e(\nu) &= \mathcal{Q}_{\nu_k, \nu_{k+1}}(x_k, x_{k+1}) e(\nu), \\ (\psi_k x_l - x_{s_k(l)} \psi_k) e(\nu) &= \begin{cases} -e(\nu) & \text{if } l = k \text{ and } \nu_k = \nu_{k+1}, \\ e(\nu) & \text{if } l = k+1 \text{ and } \nu_k = \nu_{k+1}, \\ 0 & \text{otherwise,} \end{cases} \\ (\psi_{k+1} \psi_k \psi_{k+1} - \psi_k \psi_{k+1} \psi_k) e(\nu) &= \begin{cases} \frac{\mathcal{Q}_{\nu_k, \nu_{k+1}}(x_k, x_{k+1}) - \mathcal{Q}_{\nu_k, \nu_{k+1}}(x_{k+2}, x_{k+1})}{x_k - x_{k+2}} e(\nu) & \text{if } \nu_k = \nu_{k+2}, \\ 0 & \text{otherwise,} \end{cases} \\ x_1^{\langle h_{\nu_1}, \Lambda \rangle} e(\nu) &= 0. \end{aligned}$$

The \mathbb{Z} -grading on $R^\Lambda(n)$ is given as follows:

$$\deg(e(\nu)) = 0, \quad \deg(x_k e(\nu)) = (\alpha_{\nu_k} | \alpha_{\nu_k}), \quad \deg(\psi_l e(\nu)) = -(\alpha_{\nu_l} | \alpha_{\nu_{l+1}}).$$

The following statement was proved in a special case [4, Lem.2.1]. As we are not able to find a reference for general case, we add a proof, but it is straightforward.

Lemma 2.2. *The algebra $R^\Lambda(n)$ is a finite dimensional algebra and x_1, \dots, x_n are nilpotent.*

Proof. It is proved in [17, Cor.4.4] that $R^\Lambda(n)$ is a finite dimensional algebra. Thus, we prove that x_1, \dots, x_n are nilpotent elements. Note that, by general theory of graded Artin algebras, irreducible $R^\Lambda(n)$ -modules are gradable, and it implies that any element of positive degree acts nilpotently on finite length modules. In particular, elementary symmetric functions $e_1(x_1, \dots, x_n), \dots, e_n(x_1, \dots, x_n)$ act nilpotently on $R^\Lambda(n)$. Suppose that $(\lambda_1, \dots, \lambda_n) \in \mathbf{k}^n$ is a simultaneous eigenvalue of (x_1, \dots, x_n) . Then, we have

$$e_1(\lambda_1, \dots, \lambda_n) = 0, \dots, e_n(\lambda_1, \dots, \lambda_n) = 0.$$

It follows that all λ_i are zero, implying that x_1, \dots, x_n are nilpotent. \square

For $\beta \in \mathbb{Q}^+$ with $|\beta| = n$, let

$$I^\beta = \{\nu = (\nu_1, \dots, \nu_n) \in I^n \mid \alpha_{\nu_1} + \dots + \alpha_{\nu_n} = \beta\}.$$

The symmetric group action preserves I^β , so that

$$e(\beta) = \sum_{\nu \in I^\beta} e(\nu)$$

is a central idempotent of $R^\Lambda(n)$. We define

$$R^\Lambda(\beta) = R^\Lambda(n) e(\beta).$$

The following algebras are the object of study in this paper. We do not know whether $R^\Lambda(\beta)$ is an indecomposable \mathbf{k} -algebra, for every β .

Definition 2.3. We call the algebras $R^{\Lambda_0}(\beta)$ *finite quiver Hecke algebras of type $A_{2\ell}^{(2)}$* .

2.2. Categorification of integrable modules. Let $\mathfrak{g} = \mathfrak{g}(\mathbf{A})$ be the affine Kac-Moody Lie algebra associated with the Cartan matrix \mathbf{A} . Let q be an indeterminate and we denote the corresponding quantum affine algebra by $U_q(\mathfrak{g})$. It is a $\mathbb{Q}(q)$ -algebra defined by generators e_i, f_i ($i \in I$), q^h ($h \in \mathbf{P}^\vee$) and their relations.

For $i \in I$, let $q_i = q^{d_i}$ and

$$[n]_i = \frac{q_i^n - q_i^{-n}}{q_i - q_i^{-1}}, \quad [n]_i! = \prod_{k=1}^n [k]_i.$$

For $\mathbb{A} = \mathbb{Z}[q, q^{-1}]$, we consider the \mathbb{A} -subalgebra of $U_q(\mathfrak{g})$ generated by $f_i^{(k)} := f_i^k / [k]_i!$, for $i \in I$ and $k \in \mathbb{Z}_{\geq 0}$. We denote it by $U_{\mathbb{A}}^-(\mathfrak{g})$.

Let $\Lambda \in \mathbf{P}^+$ be a dominant integral weight. Then we may consider the irreducible highest weight $U_q(\mathfrak{g})$ -module $V_q(\Lambda)$ with highest weight Λ and its \mathbb{A} -form $V_{\mathbb{A}}(\Lambda)$, which is the $U_{\mathbb{A}}^-(\mathfrak{g})$ -submodule of $V_q(\Lambda)$ generated by a fixed highest weight vector. If we specialize $V_{\mathbb{A}}(\Lambda)$ at $q = 1$, we have the irreducible highest weight \mathfrak{g} -module $V(\Lambda)$.

We denote the direct sum of split Grothendieck groups of additive categories $R^\Lambda(\beta)\text{-proj}^{\mathbb{Z}}$ of finitely generated projective graded left $R^\Lambda(\beta)$ -modules, for $\beta \in \mathbf{Q}^+$, by

$$K_0^{\mathbb{Z}}(R^\Lambda) = \bigoplus_{\beta \in \mathbf{Q}^+} K_0(R^\Lambda(\beta)\text{-proj}^{\mathbb{Z}}).$$

It has an \mathbb{A} -module structure induced by the \mathbb{Z} -grading on $R^\Lambda(n)$.

Let $e(\nu, \nu')$ be the idempotent corresponding to the concatenation $\nu * \nu'$ of ν and ν' . If $\nu' = i$, we write $e(\nu, i)$. For $\beta \in \mathbf{Q}^+$ and $i \in I$, we define functors

$$\begin{aligned} E_i : R^\Lambda(\beta + \alpha_i)\text{-mod}^{\mathbb{Z}} &\longrightarrow R^\Lambda(\beta)\text{-mod}^{\mathbb{Z}}, \\ F_i : R^\Lambda(\beta)\text{-mod}^{\mathbb{Z}} &\longrightarrow R^\Lambda(\beta + \alpha_i)\text{-mod}^{\mathbb{Z}}, \end{aligned}$$

between categories $R^\Lambda(\beta)\text{-mod}^{\mathbb{Z}}$ and $R^\Lambda(\beta + \alpha_i)\text{-mod}^{\mathbb{Z}}$ of finitely generated graded modules by $E_i(N) = e(\beta, i)N$ and $F_i(M) = R^\Lambda(\beta + \alpha_i)e(\beta, i) \otimes_{R^\Lambda(\beta)} M$, for $M \in R^\Lambda(\beta)\text{-mod}^{\mathbb{Z}}$ and $N \in R^\Lambda(\beta + \alpha_i)\text{-mod}^{\mathbb{Z}}$, respectively. Then, E_i and F_i are exact functors [17, Thm.4.5] which send projective modules to projective modules, and the following theorem holds.

Theorem 2.4 ([17, Thm.5.2]). *Set $l_i = \langle h_i, \Lambda - \beta \rangle$, for $i \in I$. Then one of the following isomorphisms of endofunctors on the category of finitely generated graded $R^\Lambda(\beta)$ -modules holds.*

(1) *If $l_i \geq 0$, then*

$$q_i^{-2} F_i E_i \oplus \bigoplus_{k=0}^{l_i-1} q_i^{2k} \text{id} \xrightarrow{\sim} E_i F_i.$$

(2) If $l_i \leq 0$, then

$$q_i^{-2} F_i E_i \xrightarrow{\sim} E_i F_i \oplus \bigoplus_{k=0}^{-l_i-1} q_i^{-2k-2} \text{id}.$$

Thus, functors $q_i^{1-\langle h_i, \Lambda - \beta \rangle} E_i$ and F_i , for $\beta \in Q^+$ and $i \in I$, define a $U_{\mathbb{A}}(\mathfrak{g})$ -module structure on $K_0^{\mathbb{Z}}(R^{\Lambda})$.

The following theorem is the cyclotomic categorification theorem conjectured by Khovanov and Lauda, and proved by Kang and Kashiwara.

Theorem 2.5 ([17, Thm.6.2]). *There exists a $U_{\mathbb{A}}(\mathfrak{g})$ -module isomorphism between $K_0^{\mathbb{Z}}(R^{\Lambda})$ and $V_{\mathbb{A}}(\Lambda)$.*

In the study of representation type, we do not need grading. So we specialize $q \rightarrow 1$. Let $R^{\Lambda}(\beta)\text{-proj}$ be the category of finitely generated projective $R^{\Lambda}(\beta)$ -modules, and define

$$K_0(R^{\Lambda}) = \bigoplus_{\beta \in Q^+} K_0(R^{\Lambda}(\beta)\text{-proj}).$$

Then we have an isomorphism of $\mathfrak{g}_{\mathbb{Z}}$ -modules $K_0(R^{\Lambda}) \cong V(\Lambda)_{\mathbb{Z}}$, where $\mathfrak{g}_{\mathbb{Z}}$ and $V(\Lambda)_{\mathbb{Z}}$ are the Kostant \mathbb{Z} -form of \mathfrak{g} and $V(\Lambda)$, respectively.

2.3. Categorification of crystals. We now recall a combinatorial realization of the crystal of $V_q(\Lambda_0)$ by *Young walls* [11, 16]. A Young wall is a generalization of a colored Young diagram based on certain level 1 perfect crystal, which gives a combinatorial realization of crystals for basic representations of various quantum affine algebras. This realization will be used in Proposition 2.8 for counting irreducible modules over finite quiver Hecke algebras. We refer the reader to [11, 16, 22, 23] for the details. For type $A_{2\ell}^{(2)}$, we consider the following blocks:

$\boxed{0}$: unit width and half-unit height, unit thickness

\boxed{i} ($i = 1, \dots, \ell$) : unit width and unit height, unit thickness

and define a *Young wall* to be a wall consisting of the above colored blocks stacked by the following rules. We write (2) for completeness, but it is a vacant condition, as we do not have blocks of half-unit thickness.

- (1) Colored blocks should be stacked in the pattern given below.
- (2) No block can be placed on top of a column of half-unit thickness.
- (3) Except for the rightmost column, there should be no free space to the right of any block.

The pattern is given as follows:

| | | | | | |
|----------|----------|----------|----------|----------|----------|
| | | | | | |
| 1 | 1 | 1 | 1 | 1 | 1 |
| 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 |
| 1 | 1 | 1 | 1 | 1 | 1 |
| \vdots | \vdots | \vdots | \vdots | \vdots | \vdots |
| ℓ | ℓ | ℓ | ℓ | ℓ | ℓ |
| \vdots | \vdots | \vdots | \vdots | \vdots | \vdots |
| 1 | 1 | 1 | 1 | 1 | 1 |
| 0 | 0 | 0 | 0 | 0 | 0 |
| | | | | | |

The sequence $(0, 1, 2, \dots, \ell - 1, \ell, \ell - 1, \dots, 2, 1, 0)$ of colors is repeated in each column. For a Young wall Y , define

$$\text{wt}(Y) = \Lambda_0 - \sum_{i \in I} k_i \alpha_i,$$

where k_i is the number of i -blocks in Y , for $i \in I$.

A column is called a *full column* if its height is a multiple of the unit length and its top is of unit thickness. As the ground state blocks have half-unit height, a column is not full if and only if the number of stacked blocks is divisible by $|\delta| = 2\ell + 1$. A Young wall is said to be *proper* if none of the full columns have the same height. A part of a column consisting of two 0-blocks, two 1-blocks, \dots , two $(\ell - 1)$ -blocks and one ℓ -block is called a δ -column. A column in a proper Young wall is said to contain a *removable* δ if we may remove a δ -column from Y and still obtain a proper Young wall.

Let $\mathcal{Y}(\Lambda_0)$ be the set of all proper Young walls Y such that none of the columns contain a removable δ . Then, Kashiwara operators \tilde{e}_i and \tilde{f}_i on $\mathcal{Y}(\Lambda_0)$ can be defined by using the combinatorics of Young walls, and $\mathcal{Y}(\Lambda_0)$ has a $U_q(\mathfrak{g})$ -crystal structure [11, 16].

Theorem 2.6. [16, Thm. 11.41] *The crystal $\mathcal{Y}(\Lambda_0)$ is isomorphic to the crystal $B(\Lambda_0)$ of the highest weight $U_q(\mathfrak{g})$ -module $V_q(\Lambda_0)$.*

Let $\mathbb{B}(\Lambda)_{\Lambda - \beta}$ be the set of isomorphism classes of irreducible $R^\Lambda(\beta)$ -modules, for $\beta \in \mathbb{Q}^+$, and define $\mathbb{B}(\Lambda) = \sqcup_{\beta \in \mathbb{Q}^+} \mathbb{B}(\Lambda)_{\Lambda - \beta}$. Then, Lauda and Vazirani defined a crystal structure on $\mathbb{B}(\Lambda)$ and proved the theorem below. We note that the idea to use socle and cosocle for defining crystal structure goes back to Leclerc's interpretation of the modular branching rule as a crystal in the Hecke algebra case.

Theorem 2.7 ([33, Thm.7.5]). *The crystal $\mathbb{B}(\Lambda)$ is isomorphic to the crystal $B(\Lambda)$ of the highest weight module $V_q(\Lambda)$.*

A realization of the crystal $B(\Lambda_0)$ in terms of partitions is easier to handle for those who are not familiar with Young walls, and we explain the realization by h -restricted h -strict partitions from [3, Th.9.2].

For $h \in \mathbb{Z}_{\geq 0}$, a partition $\lambda = (\lambda_1, \lambda_2, \dots)$ is h -strict if h divides λ_r , whenever $\lambda_r = \lambda_{r+1}$ for $r \geq 1$. If $h = 0$, we understand it as a strict partition. An h -strict partition λ is h -restricted if it satisfies $\lambda_r - \lambda_{r+1} < h$ if $h | \lambda_r$ and $\lambda_r - \lambda_{r+1} \leq h$ if $h \nmid \lambda_r$. We denote by \mathcal{RP}_h the set of all h -restricted h -strict partitions. Let

$$(2.3.1) \quad h = 2\ell + 1.$$

For a Young wall $Y \in \mathcal{Y}(\Lambda_0)$, let $\lambda_k(Y)$ be the number of stacked blocks in the k th column of Y . Then, it is easy to see that the map $Y \mapsto \lambda_Y = (\lambda_1(Y), \lambda_2(Y), \dots)$ is a 1-1 correspondence between $\mathcal{Y}(\Lambda_0)$ and \mathcal{RP}_h . Thus, \mathcal{RP}_h has the induced crystal structure, which is easy to describe, and the weight of $\lambda \in \mathcal{RP}_h$ is given by the residue pattern $\nu_\delta = 01 \dots \ell \dots 10$ on the nodes of λ . We define, for $\beta = \sum_{i \in I} k_i \alpha_i \in \mathbb{Q}^+$,

$$\mathcal{RP}_h(\beta) = \{\lambda \in \mathcal{RP}_h \mid \text{the number of } i\text{-nodes is } k_i, \text{ for } i \in I.\}$$

Then, we have the following proposition by Theorem 2.6 and Theorem 2.7.

Proposition 2.8. *For $\beta \in \mathbb{Q}^+$, the number of isomorphism classes of irreducible $R^{\Lambda_0}(\beta)$ -modules is equal to $|\mathcal{RP}_h(\beta)|$.*

3. DIMENSION FORMULA FOR $R^{\Lambda_0}(\beta)$

3.1. Fock space revisited. We need some computation in \mathcal{F} before giving $\dim R^{\Lambda_0}(\beta)$. As in the Fock space theory for type $A_{e-1}^{(1)}$, we may describe the action of Chevalley generators by the combinatorics of adding/removing nodes of Young diagrams, after minor modification. Let $\lambda \vdash n$ be a shifted Young diagram.

- Suppose that we may remove a box of residue $i \in I$ from λ and obtain a new shifted Young diagram. Then, we denote the resulting shifted Young diagram by $\lambda \nearrow \boxed{i}$.
- Similarly, if we may add a box of residue $i \in I$ to λ and obtain a new shifted Young diagram then we denote the resulting shifted Young diagram by $\lambda \swarrow \boxed{i}$.

We have $\text{wt}(\lambda \nearrow \boxed{i}) = \text{wt}(\lambda) + \alpha_i$ and $\text{wt}(\lambda \swarrow \boxed{i}) = \text{wt}(\lambda) - \alpha_i$. Recalling the residue pattern ν_δ , a box to be removed (resp. added) has residue $i \in I$ if and only if it is the rightmost box of a row of length $j + 1$ in λ (resp. $\lambda \swarrow \boxed{i}$) with $j \equiv i, -i - 1$ modulo h .

Thus, it follows from (1.3.1) and (1.3.2) that

$$(3.1.1) \quad e_i|\lambda\rangle = \sum_{\mu=\lambda \nearrow \boxed{i}} m(\lambda, \mu)|\mu\rangle, \quad f_i|\lambda\rangle = \sum_{\mu=\lambda \searrow \boxed{i}} |\mu\rangle,$$

$$\text{where } m(\lambda, \mu) = \begin{cases} 2 & \text{if } \text{wt}(\mu) = \text{wt}(\lambda) + \alpha_0 \text{ and } l(\lambda) = l(\mu), \\ 1 & \text{otherwise.} \end{cases}$$

Definition 3.1. For $\lambda \vdash n$ and $\nu \in I^n$, we denote by $K(\lambda, \nu)$ the number of standard tableaux of shape λ that has the residue sequence ν . Namely, we define

$$K(\lambda, \nu) = |\{T \in \text{ST}(\lambda) \mid \nu = \text{res}(T)\}|.$$

We have the following equality.

$$(3.1.2) \quad |\text{ST}(\lambda)| = \sum_{\nu \in I^{\Lambda_0 - \text{wt}(\lambda)}} K(\lambda, \nu).$$

In the next lemma, d is the scaling element. Note that if $\text{wt}(\lambda) = \Lambda_0 - \sum_{i \in I} k_i \alpha_i$ then $k_0 = -\langle d, \text{wt}(\lambda) \rangle$.

Lemma 3.2. Let $\lambda \vdash n$ and $\nu = (\nu_1, \dots, \nu_n) \in I^n$. Then we have

$$\begin{aligned} e_{\nu_1} e_{\nu_2} \cdots e_{\nu_n} |\lambda\rangle &= 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} K(\lambda, \nu) |0\rangle, \\ f_{\nu_n} f_{\nu_{n-1}} \cdots f_{\nu_1} |0\rangle &= \sum_{\mu \vdash n} K(\mu, \nu) |\mu\rangle. \end{aligned}$$

Proof. The second equality is clear. We prove the first equality by induction on n . Since it is obvious when $n = 1$, we assume $n > 1$. By definition, if $\mu = \lambda \nearrow \boxed{i}$, then we have

$$2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} = m(\lambda, \mu) 2^{-\langle d, \text{wt}(\mu) \rangle - l(\mu)}.$$

Let $j = \nu_n$ and $\nu^- = (\nu_1, \dots, \nu_{n-1})$. Using the induction hypothesis and (3.1.1), we obtain

$$\begin{aligned} e_{\nu_1} e_{\nu_2} \cdots e_{\nu_n} |\lambda\rangle &= \sum_{\mu=\lambda \nearrow \boxed{j}} m(\lambda, \mu) e_{\nu_1} e_{\nu_2} \cdots e_{\nu_{n-1}} |\mu\rangle \\ &= \sum_{\mu=\lambda \nearrow \boxed{j}} m(\lambda, \mu) 2^{-\langle d, \text{wt}(\mu) \rangle - l(\mu)} K(\mu, \nu^-) |0\rangle \\ &= 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} \sum_{\mu=\lambda \nearrow \boxed{j}} K(\mu, \nu^-) |0\rangle \\ &= 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} K(\lambda, \nu) |0\rangle. \end{aligned}$$

Hence the desired formula follows. \square

3.2. Dimension formula. Now we are ready to state and prove the main theorem of this section. We remark that dimension of cyclotomic quiver Hecke algebras is not known except for block algebras of cyclotomic Hecke algebras. In the case of cyclotomic Hecke algebras, we know from the theory of cellular algebras that

$$\dim R^\Lambda(\beta) = \sum_{\text{wt}(\lambda)=\Lambda-\beta} |\text{ST}(\lambda)|^2.$$

Theorem 3.4 below gives the dimension of the finite quiver Hecke algebra $R^{\Lambda_0}(\beta)$ of type $A_{2\ell}^{(2)}$ in terms of shifted standard tableaux. The idea to obtain the formula is different, and we use computation in the Fock space \mathcal{F} and the biadjointness result from [17] and [24].

We recall functors E_i and F_i from Section 2.2 and consider them in non-graded setting.

$$\begin{aligned} E_i : R^\Lambda(\beta + \alpha_i)\text{-mod} &\longrightarrow R^\Lambda(\beta)\text{-mod}, \\ F_i : R^\Lambda(\beta)\text{-mod} &\longrightarrow R^\Lambda(\beta + \alpha_i)\text{-mod}, \end{aligned}$$

Then the pair (E_i, F_i) is a biadjoint pair of exact functors by [17, Thm.4.5], [24, 3.2, Thm.3.5]. [24, 3.2] gives the adjunction $\text{id} \rightarrow E_i F_i$, $F_i E_i \rightarrow \text{id}$ on the one hand, [24, Thm.3.5] gives the adjunction $E_i F_i \rightarrow \text{id}$, $\text{id} \rightarrow F_i E_i$ on the other hand.

Lemma 3.3. *We have the following equality, for $M, N \in R^{\Lambda_0}(\beta)\text{-mod}$.*

$$\dim \text{Hom}_{R^{\Lambda_0}(\beta)}(M, N) = \dim \text{Hom}_{R^{\Lambda_0}(\beta)}(N, M).$$

Proof. We prove the assertion by induction on $|\beta|$. Suppose that the assertion holds for all $\beta' \in \mathbf{Q}^+$ such that $|\beta'| < |\beta|$. Then, by the induction hypothesis and the biadjointness of E_i and F_i , we have

$$\begin{aligned} \dim \text{Hom}_{R^{\Lambda_0}(\beta)}(F_i M, N) &= \dim \text{Hom}_{R^{\Lambda_0}(\beta - \alpha_i)}(M, E_i N) \\ &= \dim \text{Hom}_{R^{\Lambda_0}(\beta - \alpha_i)}(E_i N, M) = \dim \text{Hom}_{R^{\Lambda_0}(\beta)}(N, F_i M). \end{aligned}$$

As $R^{\Lambda_0}(\beta)$'s categorify $V(\Lambda_0)$ and

$$V(\Lambda_0)_{\Lambda_0 - \beta} = \sum_{i \in I} f_i V(\Lambda_0)_{\Lambda_0 - \beta + \alpha_i},$$

where $V(\Lambda_0)_\mu$ is the μ -weight space, the assertion follows. \square

Theorem 3.4. *Let λ be a shifted partition, $\nu \in I^n$. Then, for $\beta \in \mathbf{Q}^+$ and $\nu, \nu' \in I^\beta$, we have*

$$\begin{aligned} \dim e(\nu') R^{\Lambda_0}(n) e(\nu) &= \sum_{\lambda \vdash n} 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} K(\lambda, \nu') K(\lambda, \nu), \\ \dim R^{\Lambda_0}(\beta) &= \sum_{\lambda \vdash n, \text{wt}(\lambda) = \Lambda_0 - \beta} 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} |\text{ST}(\lambda)|^2, \\ \dim R^{\Lambda_0}(n) &= \sum_{\lambda \vdash n} 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} |\text{ST}(\lambda)|^2. \end{aligned}$$

Proof. We have $K_0(R^{\Lambda_0}) \cong V(\Lambda_0)_{\mathbb{Z}}$ by the paragraph after Theorem 2.5. Let $(\ , \)$ be the Shapovalov form on $V(\Lambda_0)$. It is a symmetric bilinear form characterized by the following properties, where v_0 is a vector which generates $\{v \in V(\Lambda_0)_{\mathbb{Z}} \mid \text{wt}(v) = \Lambda_0\}$ as a \mathbb{Z} -module.

$$(v_0, v_0) = 1, \quad (f_i x, y) = (x, e_i y), \quad \text{for } x, y \in V(\Lambda_0).$$

As (E_i, F_i) is a biadjoint pair and Lemma 3.3 holds, the above characterization implies that the induced Shapovalov form on $K_0(R^{\Lambda_0})$ is given as follows:

$$(3.2.1) \quad \dim \text{Hom}_{R^{\Lambda_0}(\beta)}(M, N) = ([M], [N]), \quad \text{for projective } R^{\Lambda_0}(\beta)\text{-modules } M \text{ and } N.$$

We identify $U(\mathfrak{g})|0\rangle = V(\Lambda_0)$ and embed $V(\Lambda_0)$ to the Fock space \mathcal{F} . As $V(\Lambda_0)$ is a direct summand of \mathcal{F} as a \mathfrak{g} -module, we may extend the Shapovalov form on $V(\Lambda_0)$ to a symmetric bilinear form on \mathcal{F} that has the property

$$(f_i x, y) = (x, e_i y), \quad \text{for } x, y \in \mathcal{F}.$$

Further, $F_i(R^{\Lambda_0}(n)e(\nu)) = R^{\Lambda_0}(n+1)e(\nu, i)$ and

$$F_{\nu_n} \dots F_{\nu_1}(R^{\Lambda_0}(0)) = R^{\Lambda_0}(n)e(\nu), \quad \text{for } \nu \in I^n.$$

Hence, for $\nu = (\nu_1, \dots, \nu_n)$, $\nu' = (\nu'_1, \dots, \nu'_n) \in I^n$, Lemma 3.2 and (3.2.1) imply that

$$\begin{aligned} \dim e(\nu') R^{\Lambda_0}(n) e(\nu) &= \dim \text{Hom}_{R^{\Lambda_0}(n)}(R^{\Lambda_0}(n)e(\nu'), R^{\Lambda_0}(n)e(\nu)) \\ &= (f_{\nu'_n} f_{\nu'_{n-1}} \dots f_{\nu'_1} |0\rangle, f_{\nu_n} f_{\nu_{n-1}} \dots f_{\nu_1} |0\rangle) \\ &= \sum_{\lambda \vdash n} K(\lambda, \nu) (f_{\nu'_n} f_{\nu'_{n-1}} \dots f_{\nu'_1} |0\rangle, |\lambda\rangle) \\ &= \sum_{\lambda \vdash n} K(\lambda, \nu) (|0\rangle, e_{\nu'_1} e_{\nu'_2} \dots e_{\nu'_n} |\lambda\rangle) \\ &= \sum_{\lambda \vdash n} 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} K(\lambda, \nu') K(\lambda, \nu). \end{aligned}$$

To deduce the second formula from the first, we use (3.1.2) and

$$R^{\Lambda_0}(\beta) = \bigoplus_{\nu, \nu' \in I^\beta} e(\nu') R^{\Lambda_0}(n) e(\nu).$$

Then we have

$$\begin{aligned} \dim R^{\Lambda_0}(\beta) &= \sum_{\nu, \nu' \in I^\beta} \sum_{\lambda \vdash n} 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} K(\lambda, \nu') K(\lambda, \nu) \\ &= \sum_{\lambda \vdash n, \text{wt}(\lambda) = \Lambda_0 - \beta} 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} \sum_{\nu \in I^\beta} \sum_{\nu' \in I^\beta} K(\lambda, \nu') K(\lambda, \nu) \\ &= \sum_{\lambda \vdash n, \text{wt}(\lambda) = \Lambda_0 - \beta} 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} |\text{ST}(\lambda)|^2. \end{aligned}$$

The last formula follows from $R^{\Lambda_0}(n) = \bigoplus_{\beta \in \mathbb{Q}^+, |\beta|=n} R^{\Lambda_0}(\beta)$. □

Theorem 3.4 has several important consequences.

Corollary 3.5. (1) Let $\nu \in I^n$. Then, $e(\nu) \neq 0$ in $R^{\Lambda_0}(n)$ if and only if ν may be obtained from a standard tableau T as $\nu = \text{res}(T)$.

(2) We have the following hook length formula, for $\beta \in \mathbb{Q}^+$.

$$\dim R^{\Lambda_0}(\beta) = \sum_{\lambda \vdash n, \text{wt}(\lambda) = \Lambda_0 - \beta} 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} \left(\frac{n!}{\prod_{(i,j) \in \lambda} h_{i,j}} \right)^2.$$

(3) For any natural number n , we have the following equality.

$$n! = \sum_{\beta \in \mathbb{Q}^+, |\beta|=n} 2^{n - \langle d, \beta \rangle} \dim R^{\Lambda_0}(\beta).$$

Proof. (1) It follows from Theorem 3.4 that

$$e(\nu) R^{\Lambda_0}(n) e(\nu) = \sum_{\lambda \vdash n} 2^{-\langle d, \text{wt}(\lambda) \rangle - l(\lambda)} K(\lambda, \nu)^2.$$

Then, observe that $e(\nu) \neq 0$ if and only if $\dim e(\nu) R^{\Lambda_0}(n) e(\nu) > 0$.

(2) It follows from Theorem 1.1(1) and Theorem 3.4.

(3) Using Theorem 1.1(2) and Theorem 3.4, we obtain

$$\begin{aligned}
n! &= \sum_{\lambda \vdash n} 2^{n-l(\lambda)} |\text{ST}(\lambda)|^2 \\
&= \sum_{\beta \in \mathbb{Q}^+, |\beta|=n} 2^{n+\langle d, \Lambda_0 - \beta \rangle} \sum_{\lambda \vdash n, |\lambda|=\beta} 2^{-\langle d, \Lambda_0 - \beta \rangle - l(\lambda)} |\text{ST}(\lambda)|^2 \\
&= \sum_{\beta \in \mathbb{Q}^+, |\beta|=n} 2^{n-\langle d, \beta \rangle} \dim R^{\Lambda_0}(\beta).
\end{aligned}$$

We have proved the equality. \square

4. REPRESENTATION TYPE OF $R^{\Lambda_0}(\beta)$

4.1. Reduction to $R^{\Lambda_0}(k\delta)$. We denote by $R^\Lambda(n)\text{-mod}$ the category of finitely generated $R^\Lambda(n)$ -modules and we define functors E and F by

$$\begin{aligned}
E &:= \sum_{i \in I} E_i : R^\Lambda(n+1)\text{-mod} \longrightarrow R^\Lambda(n)\text{-mod}, \\
F &:= \sum_{i \in I} F_i : R^\Lambda(n)\text{-mod} \longrightarrow R^\Lambda(n+1)\text{-mod}.
\end{aligned}$$

Recall that the pair (E_i, F_i) is a biadjoint pair of exact functors.

Lemma 4.1. *Let \mathcal{A} and \mathcal{B} be abelian categories, $E : \mathcal{A} \rightarrow \mathcal{B}$ an exact functor. If F is right adjoint (resp. left adjoint) to E then F sends an injective object (resp. projective object) to an injective object (resp. projective object).*

Proposition 4.2. *$R^\Lambda(\beta)$ is a self-injective algebra.*

Proof. As the pair (E_i, F_i) is a biadjoint pair of exact functors, F is right adjoint to E . Hence F preserves injective objects. On the other hand, by definition, $R^\Lambda(k+1) = FR^\Lambda(k)$ for $k \in \mathbb{Z}_{\geq 0}$. Since $R^\Lambda(0)$ is an injective $R^\Lambda(0)$ -module, $R^\Lambda(n)$ is an injective $R^\Lambda(n)$ -module and so are its direct summands $R^\Lambda(\beta)$'s. \square

In the following, we recall several general results. Let A and B be \mathbf{k} -algebras.

Theorem 4.3 (Krause). *If A and B are stable equivalent, then A and B have the same representation type.*

See [34, Thm.1] for the Krause's theorem from [31]. The following theorem is well-known.

Theorem 4.4 ([39, Thm.2.1]). *If A and B are self-injective algebras such that $D^b(A\text{-mod})$ and $D^b(B\text{-mod})$ are equivalent as triangulated categories, i.e., A and B are derived equivalent. Then A and B are stably equivalent.*

Theorem 4.5 ([5, Thm.6.4]). *Let \mathcal{A} be an artinian and noetherian \mathbf{k} -linear abelian category such that the endomorphism ring of any simple object is \mathbf{k} . Suppose that an adjoint pair (E, F) of exact endo-functors on \mathcal{A} satisfies sl_2 -categorification axioms in [5, 5.2]. Then*

- (1) *The weight space decomposition of the locally finite $sl_2(\mathbb{Q})$ -module $K(\mathcal{A}) \otimes_{\mathbb{Z}} \mathbb{Q}$ gives the decomposition of the category $\mathcal{A} = \bigoplus_{c \in \mathbb{Z}} \mathcal{A}_c$.*
- (2) *The categories \mathcal{A}_{-c} and \mathcal{A}_c are derived equivalent.*

We consider $\mathcal{A} = \bigoplus_{c \in \mathbb{Z}} \mathcal{A}_c$, where

$$\mathcal{A}_c = \bigoplus_{\mu \in \mathbf{P}: \mu(h_i) = c} R^\Lambda(\Lambda - \mu)\text{-mod.}$$

Then, (E_i, F_i) is an adjoint pair of exact endo-functors on \mathcal{A} . The following is another consequence of [24].

Theorem 4.6. *For $i \in I$, the pair of functors (E_i, F_i) satisfies the sl_2 -categorification axioms.*

Proof. It follows from [24, 3.2] that we have $X_i, X_{i+1}, T \in \text{End}(E_i^{\otimes 2})$ that satisfy nil affine Hecke relations. Thus [41, 5.3.3] implies the result. \square

Corollary 4.7. *Let Λ be an integral dominant weight, $V(\Lambda)$ the corresponding integrable highest weight \mathfrak{g} -module. If $\mu \in \mathbf{P}$ appears in its weight system, that is, if the weight space $V(\Lambda)_\mu$ is nonzero, then $R^\Lambda(\Lambda - \mu)$ and $R^\Lambda(\Lambda - w\mu)$ are derived equivalent, for $w \in \mathbf{W}$.*

Proof. As the weight system of an integrable module is \mathbf{W} -invariant, it suffices to consider the case $w = s_i$, for some $i \in I$. Then, the result follows from Theorem 4.5 and Theorem 4.6. \square

The following corollary is crucial in reducing the proof of Theorem 4.22 to the cases when β is a multiple of the null root δ .

Corollary 4.8. *Let Λ be a dominant integral weight. Then, for $w \in \mathbf{W}$ and $k \in \mathbb{Z}_{\geq 0}$, cyclotomic quiver Hecke algebras $R^\Lambda(k\delta)$ and $R^\Lambda(\Lambda - w\Lambda + k\delta)$ have the same representation type.*

Proof. Since $w\delta = \delta$, Corollary 4.7 tells that $R^\Lambda(k\delta)$ and $R^\Lambda(\Lambda - w\Lambda + k\delta)$ are derived equivalent. Then, Proposition 4.2 and Theorem 4.4 imply that they are stably equivalent. Thus, the assertion follows from Theorem 4.3. \square

In the rest of the paper, we focus on finite quiver Hecke algebras $R^{\Lambda_0}(\beta)$ of type $A_{2\ell}^{(2)}$. Then, every nonzero $R^{\Lambda_0}(\beta)$ has the form $R^{\Lambda_0}(\Lambda_0 - w\Lambda_0 + k\delta)$, which has the same representation type as $R^{\Lambda_0}(k\delta)$ by Corollary 4.8.

We recall the following theorem. Let $\text{Rad } A$ be the two-sided ideal generated by paths of positive length, for a path algebra A .

Theorem 4.9 ([9, Thm.1]). *Brauer tree algebras are stably equivalent to symmetric Nakayama algebras $A/\text{Rad}^{lm+1}A$, for the path algebra A of a cyclic quiver of length l and $m \in \mathbb{Z}_{\geq 0}$.*

Thus, Brauer tree algebras have finite representation type by Theorem 4.3. According to Professor Asashiba, arguments in Janusz [13] may be read as another proof, which is more direct and elementary. We are grateful to him for the explanation.

4.2. The algebra $R^{\Lambda_0}(\delta)$. In this subsection, we prove that $R^{\Lambda_0}(\delta)$ is a Brauer tree algebra. Hence, it has finite representation type by Theorem 4.9. We assume that $Q_{i,j}(u, v) = 1$ if $a_{ij} = 0$ for simplifying computations. This assumption is not essential. Recall that $h = 2\ell + 1$.

For $0 \leq i \leq \ell - 1$, we consider two row partitions $\lambda(i) = (h - i - 1, i) \vdash h - 1$. Note that $\text{wt}(\lambda(i)) = \Lambda_0 - \delta + \alpha_i$ and the residues are given as follows:

$$(4.2.1) \quad \begin{array}{|c|c|c|c|c|c|c|c|} \hline 0 & 1 & \cdots & i & \cdots & \ell & \cdots & i+1 \\ \hline & 0 & \cdots & i-1 & & & & \\ \hline \end{array}$$

We set

$$L_i = \bigoplus_{T \in \text{ST}(\lambda(i))} \mathbf{k}T$$

and define an $R^{\Lambda_0}(\delta - \alpha_i)$ -module structure on L_i as follows: for $\nu \in I^{\delta - \alpha_i}$, $1 \leq k \leq h - 1$ and $1 \leq l < h - 1$,

$$(4.2.2) \quad e(\nu)T = \begin{cases} T & \text{if } \nu = \text{res}(T), \\ 0 & \text{otherwise,} \end{cases} \quad x_k T = 0, \quad \psi_l T = \begin{cases} s_l T & \text{if } s_l T \text{ is standard,} \\ 0 & \text{otherwise,} \end{cases}$$

where $s_l T$ is the tableau obtained from T by exchanging the entries l and $l + 1$. We check that it is well-defined. It is easy to see from the residue pattern (4.2.1) that $\nu_k = \nu_{k+1}$ does not occur, for $\nu = \text{res}(T)$ with $T \in \text{ST}(\lambda(i))$. Thus, it is straightforward to check the defining relations except those for ψ_k^2 and for $\psi_{k+1}\psi_k\psi_{k+1} - \psi_k\psi_{k+1}\psi_k$.

Let $T \in \text{ST}(\lambda(i))$ and write $\text{res}(T) = (\nu_1, \dots, \nu_{h-1})$. Since $0 \leq i \leq \ell - 1$, one can also show from the residue pattern (4.2.1) that

- (i) Let $(1, j) \in \lambda(i)$ and $(2, j') \in \lambda(i)$ be such that $j > j'$. Then $a_{pq} = 0$, for $p = \text{res}(1, j)$ and $q = \text{res}(2, j')$.
- (ii) If $\nu_k = \nu_{k+2}$, then $(\nu_k, \nu_{k+1}, \nu_{k+2}) = (0, 1, 0)$ or $(\ell - 1, \ell, \ell - 1)$.

We show that $\psi_k^2 e(\nu) = e(\nu)$ if $a_{\nu_k, \nu_{k+1}} = 0$, and $\psi_k^2 e(\nu) = 0$ otherwise. Let $\nu = \text{res}(T)$, for $T \in \text{ST}(\lambda(i))$. Since (i) implies that $s_k T$ is standard if and only if $a_{\nu_k, \nu_{k+1}} = 0$, we have

$$\psi_k^2 T = \begin{cases} T & \text{if } a_{\nu_k, \nu_{k+1}} = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Thus, we have proved that $\psi_k^2 e(\nu) = Q_{\nu_k, \nu_{k+1}}(x_k, x_{k+1})e(\nu)$ holds on L_i .

As $s_k T, s_{k+1} s_k T, s_k s_{k+1} s_k T$ are all standard if and only if $s_{k+1} T, s_k s_{k+1} T, s_{k+1} s_k s_{k+1} T$ are all standard, the operators ψ_k and ψ_{k+1} on L_i satisfy the Artin braid relation. Thus, we show that the terms with $\nu_k = \nu_{k+2}$ in the defining relation vanish. Since (ii) implies that we only have to consider $(\nu_k, \nu_{k+1}, \nu_{k+2}) = (0, 1, 0)$ or $(\ell - 1, \ell, \ell - 1)$, and $\deg(\psi_k \psi_{k+1} \psi_k e(\nu)) > 0$ in both cases, they vanish as desired.

Lemma 4.10. *For $i = 0, 1, \dots, \ell - 1$,*

- (1) $R^{\Lambda_0}(\delta - \alpha_i)$ is a simple algebra,
- (2) L_i is an irreducible $R^{\Lambda_0}(\delta - \alpha_i)$ -module of dimension $\binom{h-2}{i} - \binom{h-2}{i-1}$.

Proof. (1) Since $\Lambda_0 - \delta + \alpha_i$ is a maximal weight of $V(\Lambda_0)$,

$$\dim V(\Lambda_0)_{\Lambda_0 - \delta + \alpha_i} = 1, \quad \Lambda_0 - \delta + \alpha_i = w\Lambda_0,$$

for some $w \in W$ [15, Lem.12.6]. Then, as $R^{\Lambda_0}(0)$ is a simple algebra, the assertion follows from Corollary 4.8 and the fact that derived equivalence preserves the center.

(2) Theorem 1.1 gives the dimension of L_i as follows:

$$\dim L_i = \frac{(h-1)!}{(h-1) \left(\frac{(h-i-1)!}{h-2i-1} \right) i!} = \frac{(h-2)!}{(h-i-2)!i!} - \frac{(h-2)!}{(h-i-1)!(i-1)!}.$$

Next we show that $\dim e(\nu)L_i \leq 1$, for any ν . Then, a standard argument shows that L_i is irreducible. But, if one tries to enumerate all possible standard tableaux T with $\text{res}(T) = \nu$, one finds that it is unique if it exists. It follows that $\dim e(\nu)L_i \leq 1$. \square

We now extend the $R^{\Lambda_0}(\delta - \alpha_i)$ -module L_i to an $R^{\Lambda_0}(\delta)$ -module S_i as follows. As the generators x_k , for $1 \leq k \leq h-1$, and ψ_l , for $1 \leq l < h-1$, act as (4.2.2), we define the action of x_h and ψ_{h-1} . We declare that both act as 0. The idempotents $e(\nu)$, for $\nu \in I^h$, act as

$$e(\nu)T = \begin{cases} T & \text{if } \nu = \text{res}(T) * i, \\ 0 & \text{otherwise.} \end{cases}$$

Here, $\text{res}(T) * i$ is the sequence obtained from $\text{res}(T)$ by adding i at the right end. We check that it is well-defined. Let $\nu = \text{res}(T) * i$. Then, it follows from (4.2.1) that

$$\nu_{h-2} \neq i \quad \text{and} \quad \nu_{h-1} = i-1 \text{ or } i+1,$$

if $\ell \geq 2$, $\deg Q_{\nu_{h-2}, \nu_{h-1}} \geq 2$ and $\nu_{h-1} = i \pm 1$ if $\ell = 1$. Then, $\nu_{h-1} = i \pm 1$ implies that $\nu_{h-1} = \nu_h$ does not occur. Thus, the relations which involve x_h all hold. It also follows that $a_{\nu_{h-1}\nu_h} = 0$ does not occur, so that the relation for ψ_{h-1}^2 holds. Finally, $\nu_{h-2} \neq \nu_h$ if $\ell \geq 2$ and $\deg Q_{\nu_{h-2}, \nu_{h-1}} \geq 2$ if $\ell = 1$ implies that the relation for $\psi_{h-1}\psi_{h-2}\psi_{h-1} - \psi_{h-2}\psi_{h-1}\psi_{h-2}$. Note that S_i is a homogeneous representation in the sense of Kleshchev and Ram [28], and

$$(4.2.3) \quad E_j S_i = \begin{cases} L_i & \text{if } j = i, \\ 0 & \text{if } j \neq i. \end{cases}$$

Lemma 4.11. *The set $\{S_i \mid 0 \leq i \leq \ell - 1\}$ is a complete set of irreducible $R^{\Lambda_0}(\delta)$ -modules.*

Proof. By Proposition 2.8, the number of pairwise non-isomorphic irreducible $R^{\Lambda_0}(\delta)$ -modules is ℓ . On the other hand, (4.2.3) tells us that S_i 's are pairwise non-isomorphic. Thus, we have the assertion. \square

We now consider the module $F_i L_i$. Note that $F_i L_i$ is a projective-injective module since the functor F_i preserves projectivity and injectivity. By the biadjointness of E_i and F_i ,

$$(4.2.4) \quad \begin{aligned} \text{Hom}(S_j, F_i L_i) &\simeq \text{Hom}(E_i S_j, L_i) \simeq \begin{cases} \mathbf{k} & \text{if } j = i, \\ 0 & \text{if } j \neq i, \end{cases} \\ \text{Hom}(F_i L_i, S_j) &\simeq \text{Hom}(L_i, E_i S_j) \simeq \begin{cases} \mathbf{k} & \text{if } j = i, \\ 0 & \text{if } j \neq i, \end{cases} \end{aligned}$$

which implies that $F_i L_i$ is indecomposable and $\text{Top}(F_i L_i) = \text{Soc}(F_i L_i) = S_i$. So, $F_i L_i$ is the projective cover of S_i .

Theorem 4.12. *If $\ell \geq 2$, the radical series of $F_i L_i$, for $0 \leq i \leq \ell - 1$, is given as follows:*

$$\begin{array}{ccccc} & S_0 & & S_i & & S_{\ell-1} \\ F_0 L_0 \simeq & S_0 \oplus S_1 & , & F_i L_i \simeq & S_{i-1} \oplus S_{i+1} & (i \neq 0, \ell - 1), & F_{\ell-1} L_{\ell-1} \simeq & S_{\ell-2} & . \\ & S_0 & & S_i & & S_{\ell-1} \end{array}$$

If $\ell = 1$, S_0 is the unique irreducible module and $F_0 L_0$ is a uniserial module of length 3.

Proof. By definition, $E_j L_i = 0$, for $j \neq i \pm 1$, and we know that

$$E_j L_i \simeq E_i L_j, \text{ for } j = i \pm 1$$

by comparing their characters, since both are irreducible. For $i \neq j$, by the biadjointness of E_j and F_j and [17, Thm.5.1], we have

$$\text{Hom}(F_i L_i, F_j L_j) \simeq \text{Hom}(E_j F_i L_i, L_j) \simeq \text{Hom}(F_i E_j L_i, L_j) \simeq \text{Hom}(E_j L_i, E_i L_j),$$

which gives

$$\dim \operatorname{Hom}(F_i L_i, F_j L_j) = \begin{cases} 1 & \text{if } j = i \pm 1, \\ 0 & \text{if } j \neq i, i \pm 1. \end{cases}$$

We now consider the case $i = j$. Since $E_i L_i = 0$ and $\langle h_i, \Lambda_0 - \delta + \alpha_i \rangle > 0$, we have, by Theorem 2.4,

$$E_i F_i L_i \simeq L_i^{\oplus \langle h_i, \Lambda_0 - \delta + \alpha_i \rangle}$$

So, we have

$$\operatorname{Hom}(F_i L_i, F_i L_i) \simeq \operatorname{Hom}(L_i, E_i F_i L_i) \simeq \mathbf{k}^{\oplus \langle h_i, \Lambda_0 - \delta + \alpha_i \rangle}.$$

To summarize, we have the following decomposition numbers.

$$[F_i L_i : S_j] = \begin{cases} 0 & \text{if } j \neq i \pm 1, \\ 1 & \text{if } j = i \pm 1, \\ 2 & \text{if } j = i \neq 0, \\ 3 & \text{if } j = i = 0. \end{cases}$$

In other words, we have

$$\begin{aligned} [F_0 L_0] &= \begin{cases} 3[S_0] + [S_1] & \text{if } \ell \geq 2, \\ 3[S_0] & \text{if } \ell = 1, \end{cases} \\ [F_i L_i] &= 2[S_i] + [S_{i-1}] + [S_{i+1}] \quad (1 \leq i \leq \ell - 2), \\ [F_{\ell-1} L_{\ell-1}] &= 2[S_{\ell-1}] + [S_{\ell-2}] \quad \text{if } \ell \geq 2, \end{aligned}$$

in the Grothendieck group $K_0(R^{\Lambda_0}(\delta)\text{-mod})$. Since $\operatorname{Top}(F_i L_i) = \operatorname{Soc}(F_i L_i) = S_i$, we only have to show that the heart of $F_i L_i$ is a direct sum of two irreducible modules when $i \neq \ell - 1$. Recall that the anti-involution of $R^{\Lambda_0}(\delta)$, which fixes the generators elementwise, defines a left module structure on the \mathbf{k} -dual of any module. Further, the character consideration shows that irreducible modules are self-dual. This implies that the \mathbf{k} -dual of $F_i L_i$ is isomorphic to $F_i L_i$ itself. It follows that the heart of $F_i L_i$ is self-dual, and the assertion follows. \square

As a corollary, we have the following theorem, for $\ell \geq 1$. (1) is by definition of the Brauer tree algebra, and (2) follows from Theorem 4.9.

Theorem 4.13. (1) $R^{\Lambda_0}(\delta)$ is the Brauer tree algebra associated with Brauer graph

$$\begin{array}{c} \bullet \xrightarrow{S_0} \circ \xrightarrow{S_1} \circ \xrightarrow{\cdots} \circ \xrightarrow{S_{\ell-2}} \circ \xrightarrow{S_{\ell-1}} \circ \cdot \\ e = 2 \end{array}$$

(2) $R^{\Lambda_0}(\delta)$ is not semisimple and has finite representation type.

4.3. The algebra $R^{\Lambda_0}(2\delta)$. We define $\nu^k \in I^{h+k}$, for $1 \leq k \leq h$, by adding residues of ν_δ one by one. Namely,

$$\begin{aligned} \nu^1 &= \nu_\delta * (0), \nu^2 = \nu_\delta * (01), \dots, \nu^{\ell+1} = \nu_\delta * (01 \cdots \ell), \dots \\ \dots, \nu^{h-1} &= \nu_\delta * (012 \cdots \ell \cdots 21), \nu^h = \nu_\delta * \nu_\delta. \end{aligned}$$

We denote the corresponding idempotent by $e_k = e(\nu^k)$. Then, $e_k \in R^{\Lambda_0}(\beta_k)$, where β_k are

$$\begin{aligned} \beta_1 &= \delta + \alpha_0, \beta_2 = \delta + \alpha_0 + \alpha_1, \dots, \beta_{\ell+1} = \delta + \alpha_0 + \cdots + \alpha_\ell, \dots \\ \dots, \beta_{h-1} &= 2\delta - \alpha_0, \beta_h = 2\delta. \end{aligned}$$

We define $i_k \in I$, for $1 \leq k < h$, by $\alpha_{i_k} = \beta_{k+1} - \beta_k$.

Lemma 4.14. (1) $\dim e_k R^{\Lambda_0}(\beta_k) e_k = 12$, for $1 \leq k \leq h-1$.

(2) We have an isomorphism of \mathbf{k} -algebras

$$e_k R^{\Lambda_0}(\beta_k) e_k \cong e_{k+1} R^{\Lambda_0}(\beta_{k+1}) e_{k+1}, \text{ for } 1 \leq k < h-1,$$

which sends $x_h e_k, x_{h+1} e_k, \psi_h e_k$ to $x_h e_{k+1}, x_{h+1} e_{k+1}, \psi_h e_{k+1}$, respectively.

Proof. (i) We use Theorem 3.4. In counting standard tableaux T with $\text{res}(T) = \nu^k$, one finds that the filling of $1, \dots, h-1$ are unique, the filling of $h, h+1$ is either $(1, h), (1, h+1)$ or $(1, h), (2, 2)$, and after that, there is unique way to fill in $h+2, \dots, h+k$. Thus,

$$K(\lambda, \nu_k) = \begin{cases} 1 & \text{if } \lambda = (h+k), \\ 2 & \text{if } \lambda = (h, k), \\ 0 & \text{otherwise.} \end{cases}$$

It follows that $\dim e_k R^{\Lambda_0}(\beta_k) e_k = 2^2 \cdot 1^2 + 2^1 \cdot 2^2 = 12$.

(ii) We can easily find that

$$\begin{aligned} \langle h_0, \Lambda_0 - \delta \rangle, \langle h_1, \Lambda_0 - \delta - \alpha_0 \rangle, \dots, \langle h_\ell, \Lambda_0 - \delta - \alpha_0 - \cdots - \alpha_{\ell-1} \rangle, \\ \langle h_{\ell-1}, \Lambda_0 - 2\delta + \alpha_0 + \cdots + \alpha_{\ell-1} \rangle, \dots, \langle h_1, \Lambda_0 - 2\delta + \alpha_0 + \alpha_1 \rangle, \end{aligned}$$

are all positive, for $\ell \geq 1$. Thus, Theorem 2.4 implies that there is a $(R^{\Lambda_0}(\beta_k), R^{\Lambda_0}(\beta_k))$ -bimodule monomorphism

$$R^{\Lambda_0}(\beta_k) \rightarrow e(\beta_k, i_k) R^{\Lambda_0}(\beta_{k+1}) e(\beta_k, i_k),$$

for $1 \leq k < h-1$, which respects generators. Thus, we have an $(e_k R^{\Lambda_0}(\beta_k) e_k, e_k R^{\Lambda_0}(\beta_k) e_k)$ -bimodule monomorphism

$$e_k R^{\Lambda_0}(\beta_k) e_k \rightarrow e_{k+1} R^{\Lambda_0}(\beta_{k+1}) e_{k+1},$$

for $1 \leq k < h-1$, such that $x_h e_k, x_{h+1} e_k, \psi_h e_k \mapsto x_h e_{k+1}, x_{h+1} e_{k+1}, \psi_h e_{k+1}$, respectively. As both algebras are 12-dimensional, it is a \mathbf{k} -algebra isomorphism. \square

Definition 4.15. The cyclotomic nilHecke algebra NH_2^m , for a positive integer m , is the \mathbf{k} -algebra defined by generators y_1, y_2, ψ and the relations

$$y_1^m = 0, \quad y_2 \psi - \psi y_1 = 1 = \psi y_2 - y_1 \psi, \quad \psi^2 = 0.$$

We need the fact $NH_2^3 \cong \text{Mat}(2, \mathbf{k}[x]/(x^3))$ [32, (5.4)] in the proof of next proposition.

Lemma 4.16. *We have $x_{h-1} e(s_{h-1} \nu^1) = 0$.*

Proof. If $\ell = 1$, then $e(s_2 \nu^1) = 0$, so that there is nothing to prove. Suppose that $\ell \geq 2$ and $Q_{i,j}(u, v) = 1$ if $a_{ij} = 0$ and

$$Q_{i,j}(u, v) = \begin{cases} u + v & \text{if } a_{ij} = a_{ji} = -1, \\ u^2 + v & \text{if } a_{ij} = -2 \text{ and } a_{ji} = -1, \end{cases}$$

for simplifying computation. This assumption is not essential.

We define μ by

$$\mu = s_3 s_4 \cdots s_{h-1} \nu^1 = (0, 1, 0, 2, 3, \dots, \ell-1, \ell, \ell-1, \dots, 1, 0).$$

By Corollary 3.5, and the defining relations, we have

- (i) $e(s_1 \mu) = e(s_2 \mu) = 0$,
- (ii) For $3 \leq r \leq h-2$, we have

$$\psi_r \psi_{r-1} \cdots \psi_3 e(\mu) \psi_3 \cdots \psi_{r-1} \psi_r = e(s_{r+1} s_{r+2} \cdots s_{h-1} \nu^1).$$

Then, it follows that

$$x_3 e(\mu) = (x_1 + x_3) e(\mu) = (\psi_2 \psi_1 \psi_2 - \psi_1 \psi_2 \psi_1) e(\mu) = 0,$$

which yields

$$\begin{aligned} x_{h-1} e(s_{h-1} \nu^1) &= x_{h-1} (\psi_{h-2} \cdots \psi_3 e(\mu) \psi_3 \cdots \psi_{h-2}) \\ &= \psi_{h-2} \cdots \psi_3 x_3 e(\mu) \psi_3 \cdots \psi_{h-2} \\ &= 0. \end{aligned}$$

We have proved the result. \square

Proposition 4.17. (1) *The following set spans $e_1 R^{\Lambda_0}(\beta_1) e_1$ as a \mathbf{k} -vector space.*

$$\left\{ x_h^a x_{h+1}^b e_1, \quad x_h^a x_{h+1}^b \psi_h e_1 \mid a, b \in \mathbb{Z}_{\geq 0} \right\}.$$

(2) We have the following isomorphism of \mathbf{k} -algebras.

$$NH_2^3 \cong e_1 R^{\Lambda_0}(\beta_1) e_1 : y_1 \mapsto x_h e_1, y_2 \mapsto x_{h+1} e_1, \psi \mapsto \psi_h e_1.$$

(3) The algebra $e_{h-1} R^{\Lambda_0}(\beta_{h-1}) e_{h-1}$ is generated by $x_h e_{h-1}, x_{h+1} e_{h-1}, \psi_h e_{h-1}$, which obey the affine nilHecke relations, and we have the following isomorphism of \mathbf{k} -algebras.

$$e_{h-1} R^{\Lambda_0}(\beta_{h-1}) e_{h-1} \cong \text{Mat}(2, \mathbf{k}[x]/(x^3)).$$

Proof. (1) In the proof, we assume that $Q_{i,j}(u, v) = 1$ if $a_{ij} = 0$ and

$$Q_{i,j}(u, v) = \begin{cases} u + v & \text{if } a_{ij} = a_{ji} = -1, \\ u^2 + v & \text{if } a_{ij} = -2 \text{ and } a_{ji} = -1, \\ u^4 + v & \text{if } a_{ij} = -4 \text{ and } a_{ji} = -1, \end{cases}$$

for simplifying computation. This assumption is not essential. Let $e = e_1$ and $\nu = \nu^1$. Observe that $s_i \nu$, for $1 \leq i \leq h-2$, can not be the residue sequence of a standard tableau. Thus, $e\psi_i = 0 = \psi_i e$, for $1 \leq i \leq h-2$, by Corollary 3.5. Suppose that $\ell \geq 2$. We have

$$e\psi_1^2 e = (x_1^2 + x_2)e, \quad e\psi_i^2 e = (x_i + x_{i+1})e, \text{ for } 2 \leq i \leq \ell-1, \quad e\psi_\ell^2 e = (x_\ell^2 + x_{\ell+1})e,$$

so that, starting with $x_1 = 0$, we obtain $x_i e = 0$, for $1 \leq i \leq \ell+1$. $x_{\ell+2} e = 0$ follows from

$$e\psi_{\ell+1}\psi_\ell\psi_{\ell+1}e - e\psi_\ell\psi_{\ell+1}\psi_\ell e = (x_\ell + x_{\ell+2})e.$$

Then, $e\psi_i^2 e = (x_i + x_{i+1})e$, for $\ell+2 \leq i \leq h-2$, proves that $x_i e = 0$, for $1 \leq i \leq h-1$. If $\ell = 1$, then $x_1 = 0$ and $e\psi_1^2 e = (x_1^2 + x_2)e$ proves $x_i e = 0$, for $1 \leq i \leq h-1$.

For each $w \in S_n$, we fix a reduced expression and defines ψ_w . Then, $e_1 R^{\Lambda_0}(\beta_1) e_1$ is spanned by the set

$$\left\{ x_h^a x_{h+1}^b e\psi_w e \mid a, b \in \mathbb{Z}_{\geq 0}, w \in S_{h+1} \right\}.$$

Hence, it is enough to show that $e\psi_w e = 0$ unless $w = 1$ or $w = s_h$. Recall the distinguished coset representatives for $S_{n-1} \backslash S_n$:

$$S_n = \bigsqcup_{i=1}^n S_{n-1} s_{n-1} \cdots s_i.$$

Hence, we may choose the reduced expression as $w = w_2 \cdots w_h$, where

$$\begin{aligned} w_h &\in \{s_h s_{h-1} \cdots s_1, s_h s_{h-1} \cdots s_2, \dots, s_h, 1\} \\ w_{h-1} &\in \{s_{h-1} s_{h-2} \cdots s_1, s_{h-1} s_{h-2} \cdots s_2, \dots, s_{h-1}, 1\} \\ &\dots\dots\dots \\ w_2 &\in \{s_1, 1\}. \end{aligned}$$

Suppose that $e\psi_w e \neq 0$. If $w_2 \neq 1$, $e\psi_1 = 0$ implies $e\psi_w e = 0$, so that $w_2 = 1$. Arguing similarly, we obtain $w_2 = \cdots = w_{h-2} = 1$. Using $\psi_i e = 0$, for $1 \leq i \leq h-2$, we deduce that

$$w_h \in \{s_h s_{h-1}, s_h, 1\}.$$

But, $s_h s_{h-1} \nu = 01 \cdots \ell \cdots 2001$ if $\ell \geq 2$, and $s_h s_{h-1} \nu = 0001$ if $\ell = 1$. In either case, it cannot be of the form $\text{res}(T)$, for a standard tableau T . Hence, we have either $w_h = 1$ or $w_h = s_h$. It follows that $e\psi_w e = e\psi_{w_{h-1}} e\psi_{w_h}$, and using $\psi_i e = 0$, for $1 \leq i \leq h-2$ again, we have either $w_{h-1} = 1$ or $w_{h-1} = s_{h-1}$. But $w_{h-1} = s_{h-1}$ implies $e\psi_w e = e(\nu)e(s_{h-1}\nu)\psi_{h-1}\psi_{w_h} = 0$ because $s_{h-1}\nu \neq \nu$, and we must have $w_{h-1} = 1$. We have proved (1).

(2) If $\ell \geq 2$, then $e_1 \psi_{h-1}^2 e_1 = (x_{h-1} + x_h^2)e_1 = x_h^2 e_1$ and Lemma 4.16 implies

$$x_h^3 e_1 = e_1 \psi_{h-1}^2 e_1 x_h = e_1 \psi_{h-1} x_{h-1} e(s_{h-1}\nu^1) \psi_{h-1} e_1 = 0.$$

If $\ell = 1$, then $e_1 \psi_i = 0 = \psi_i e_1$, for $i = 1, 2$, implies

$$x_3^3 e_1 = (x_1^3 + x_1^2 x_3 + x_1 x_3^2 + x_3^3) e_1 = e_1 (\psi_2 \psi_1 \psi_2 - \psi_1 \psi_2 \psi_1) e_1 = 0.$$

Thus, $x_h^3 e_1 = 0$, for all $\ell \geq 1$, and we may define a homomorphism of \mathbf{k} -algebras

$$NH_2^3 \rightarrow e_1 R^{\Lambda_0}(\beta_1) e_1 : y_1 \mapsto x_h e_1, y_2 \mapsto x_{h+1} e_1, \psi \mapsto \psi_h e_1,$$

which is an epimorphism by (1). As both algebras are 12 dimensional, it is an isomorphism.

(3) follows from (2) and Lemma 4.14(2). \square

Proposition 4.18. *The algebra $e_h R^{\Lambda_0}(2\delta) e_h$ is a quotient algebra of $\text{Mat}(2, \mathbf{k}[x]/(x^3)) \otimes_{\mathbf{k}} \mathbf{k}[t]$ by the ideal generated by an element of the form $t^3 - c_2 t^2 - c_1 t - c_0$, for some*

$$c_0, c_1, c_2 \in \text{Mat}(2, \mathbf{k}[x]/(x^3)).$$

Proof. By explicit enumeration of possible standard tableaux, Theorem 3.4 computes

$$\dim e_h R^{\Lambda_0}(2\delta) e_h = 36.$$

We consider the algebra homomorphism

$$e_{h-1} R^{\Lambda_0}(\beta_{h-1}) e_{h-1} \otimes_{\mathbf{k}} \mathbf{k}[t] \longrightarrow e_h R^{\Lambda_0}(2\delta) e_h$$

defined by $m \otimes t^k \mapsto m x_{2h}^k e_h$, for $m \in e_{h-1} R^{\Lambda_0}(\beta_{h-1}) e_{h-1}$ and $k \in \mathbb{Z}_{\geq 0}$.

This algebra homomorphism is well-defined because x_{2h} commutes with $x_h e_h, x_{h+1} e_h, \psi_h e_h$, and $x_h e_{h-1}, x_{h+1} e_{h-1}, \psi_h e_{h-1}$ generate $e_{h-1} R^{\Lambda_0}(\beta_{h-1}) e_{h-1}$ by Proposition 4.17(3).

As $\langle h_0, \Lambda_0 - 2\delta + \alpha_0 \rangle = 3$, Theorem 2.4 implies that its restriction to

$$\bigoplus_{k=0,1,2} e_{h-1} R^{\Lambda_0}(\beta_{h-1}) e_{h-1} \otimes t^k \longrightarrow e_h R^{\Lambda_0}(2\delta) e_h$$

is an isomorphism of \mathbf{k} -vector spaces. In fact, the isomorphism of functors in Theorem 2.4 is given in an explicit manner, and the terms in the direct sum are given by multiplication by x_{2h}^k , for $0 \leq k \leq 2$. See the statement of [24, Thm.3.4], for example.

Thus, $t^3 = c_2 t^2 + c_1 t + c_0$, for some $c_0, c_1, c_2 \in e_{h-1} R^{\Lambda_0}(\beta_{h-1}) e_{h-1}$, and if we consider the factor algebra by the cubic relation, the algebra homomorphism induces an isomorphism to $e_h R^{\Lambda_0}(2\delta) e_h$. Recalling the isomorphism of \mathbf{k} -algebras

$$e_{h-1} R^{\Lambda_0}(\beta_{h-1}) e_{h-1} \cong \text{Mat}(2, \mathbf{k}[x]/(x^3))$$

from Proposition 4.17(3), we have the result. \square

Proposition 4.19. *The algebra $R^{\Lambda_0}(2\delta)$ is wild, for all $\ell \geq 1$.*

Proof. Let $A = e_h R^{\Lambda_0}(2\delta) e_h$. It suffices to show that A is wild. We define a two-sided ideal of $\text{Mat}(2, \mathbf{k}[x]/(x^3)) \otimes \mathbf{k}[t]$ by

$$J = \text{Mat}(2, \mathbf{k}[x]/(x^3)) \otimes t\mathbf{k}[t] + \text{Mat}(2, x\mathbf{k}[x]/(x^3)) \otimes \mathbf{k}[t].$$

We know from Proposition 4.18 that $A/\text{Rad}^3 A$ is isomorphic to

$$\text{Mat}(2, \mathbf{k}[x]/(x^3)) \otimes \mathbf{k}[t]/J^3$$

as \mathbf{k} -algebras. We show that this algebra is wild. Then it follows that A is wild. Let E_{11} be the matrix unit. Then, for the idempotent $f = E_{11} \otimes 1$, we have

$$f (\text{Mat}(2, \mathbf{k}[x]/(x^3)) \otimes \mathbf{k}[t]/J^3) f \cong \mathbf{k}[x, t]/(x^3, x^2 t, x t^2, t^3).$$

In [40, (1.2)], the following result is attributed to Brenner and Drozd:

the commutative algebra $\mathbf{k}[x, t]/(x^2, x t^2, t^3)$ is wild.

As $\mathbf{k}[x, t]/(x^2, x t^2, t^3)$ is a quotient algebra of $\mathbf{k}[x, t]/(x^3, x^2 t, x t^2, t^3)$, $\mathbf{k}[x, t]/(x^3, x^2 t, x t^2, t^3)$ is wild. We have proved that A is wild. \square

4.4. Representation type of $R^{\Lambda_0}(\beta)$. It remains to consider $R^{\Lambda_0}(k\delta)$, for $k \geq 3$.

Lemma 4.20 ([8, Prop.2.3]). *Let A and B be finite dimensional \mathbf{k} -algebras and suppose that there exist a constant $C > 0$ and functors*

$$F : A\text{-mod} \rightarrow B\text{-mod}, \quad G : B\text{-mod} \rightarrow A\text{-mod}$$

such that, for any A -module M ,

- (1) *M is a direct summand of $GF(M)$ as an A -module,*
- (2) *$\dim F(M) \leq C \dim M$.*

If A is wild then so is B .

Proposition 4.21. *$R^{\Lambda_0}(k\delta)$, for $k \geq 2$, are wild.*

Proof. Suppose that $\ell = 1$. Then the numbers

$$\langle h_0, \Lambda_0 - k\delta \rangle = 1, \quad \langle h_1, \Lambda_0 - k\delta - \alpha_0 \rangle = 1, \quad \langle h_0, \Lambda_0 - k\delta - \alpha_0 - \alpha_1 \rangle = 3,$$

are all positive. Thus, the functors

$$\begin{aligned} F_0 &: R^{\Lambda_0}(k\delta)\text{-mod} \rightarrow R^{\Lambda_0}(k\delta + \alpha_0)\text{-mod}, \\ F_1 &: R^{\Lambda_0}(k\delta + \alpha_0)\text{-mod} \rightarrow R^{\Lambda_0}(k\delta + \alpha_0 + \alpha_1)\text{-mod}, \\ F_0 &: R^{\Lambda_0}(k\delta + \alpha_0 + \alpha_1)\text{-mod} \rightarrow R^{\Lambda_0}((k+1)\delta)\text{-mod} \end{aligned}$$

satisfy the assumptions of Lemma 4.20, by Theorem 2.4. As $R^{\Lambda_0}(2\delta)$ is wild by Proposition 4.19, so are $R^{\Lambda_0}(k\delta)$, for $k \geq 3$. Suppose that $\ell \geq 2$. Then

$$\begin{aligned} \langle h_0, \Lambda_0 - k\delta \rangle, \dots, \langle h_\ell, \Lambda_0 - k\delta - \alpha_0 - \dots - \alpha_{\ell-1} \rangle, \dots \\ \dots, \langle h_1, \Lambda_0 - (k+1)\delta + \alpha_0 + \alpha_1 \rangle, \dots, \langle h_0, \Lambda_0 - (k+1)\delta + \alpha_0 \rangle, \end{aligned}$$

are all positive, and the same argument as $\ell = 1$ case proves the result. \square

We recall that $V(\Lambda_0)_\mu \neq 0$ if and only if $\mu = \kappa - k\delta$, for some $\kappa \in W\Lambda_0$ and $k \in \mathbb{Z}_{\geq 0}$, and the pair $(\kappa, k) \in W\Lambda_0 \times \mathbb{Z}_{\geq 0}$ is uniquely determined by μ . The following theorem is the Erdmann-Nakano type theorem for type $A_{2\ell}^{(2)}$.

Theorem 4.22. *Let $\kappa \in W\Lambda_0$ and $k \in \mathbb{Z}_{\geq 0}$. The finite quiver Hecke algebra $R^{\Lambda_0}(\Lambda_0 - \kappa + k\delta)$ of type $A_{2\ell}^{(2)}$, for $\ell \geq 1$, is*

- (1) *simple if $k = 0$,*
- (2) *of finite representation type but not semisimple if $k = 1$,*
- (3) *of wild representation type if $k \geq 2$,*
- (4) *and tame representation type does not occur.*

Proof. As the representation type of $R^{\Lambda_0}(\Lambda_0 - \kappa + k\delta)$ is the same as that of $R^{\Lambda_0}(k\delta)$ by Corollary 4.8, the result follows from Theorem 4.13 and Proposition 4.21. \square

5. APPENDIX: GENERALIZED CELLULAR STRUCTURE

5.1. Generalized cellular algebras. We first recall König and Xi's notion of *affine cellular algebra* [30]. Let \mathbf{k} be a field. We note that they also require that B is commutative in the definition below.

Definition 5.1. Let (A, σ_A) and (B, σ_B) be \mathbf{k} -algebras with anti-involution. A two-sided ideal $J \subseteq A$ is called an *affine cell ideal* if $\sigma_A(J) = J$ and there exist an (A, B) -bimodule C and an (A, A) -bimodule isomorphism

$$\alpha : J \simeq C \otimes_B C^{\text{op}},$$

where C^{op} is C equipped with right A -module structure given by $xa = \sigma_A(a)x$, for $x \in C$ and $a \in A$, such that

- (a) C is free as a B -module.
- (b) Let $\tau_A : C \otimes_B C^{\text{op}} \rightarrow C \otimes_B C^{\text{op}}$ be the flip $x \otimes y \mapsto y \otimes x$. Then $\alpha \circ \sigma_A = \tau_A \circ \alpha$.

Recently, Masaki Mori has modified the definition and introduced more transparent general setup.

Definition 5.2. Let A and B be \mathbf{k} -algebras. We call a pair of (A, B) -bimodule M and (B, A) -bimodule N *weak Morita pair* if we have (A, A) and (B, B) -bimodule homomorphisms

$$\varphi_A : M \otimes_B N \longrightarrow A \quad \text{and} \quad \varphi_B : N \otimes_A M \longrightarrow B$$

such that the following diagrams commute:

$$\begin{array}{ccc} M \otimes_B N \otimes_A M & \xrightarrow{\varphi_A \otimes \text{id}} & A \otimes_A M \\ \text{id} \otimes \varphi_B \downarrow & & \downarrow \wr \\ M \otimes_B B & \xrightarrow{\sim} & M, \end{array} \quad \begin{array}{ccc} N \otimes_A M \otimes_B N & \xrightarrow{\varphi_B \otimes \text{id}} & B \otimes_B N \\ \text{id} \otimes \varphi_A \downarrow & & \downarrow \wr \\ N \otimes_A A & \xrightarrow{\sim} & N. \end{array}$$

Suppose that (M, N) is a weak Morita pair. We denote $J_A = \text{Im}(\varphi_A)$ and $J_B = \text{Im}(\varphi_B)$. Mori has proved the following lemma by showing that any element in $M \otimes_B V \setminus \text{Ker}(\Phi)$ generates $M \otimes_B V$: the proof resembles the proof of [36, Prop.2.11].

Lemma 5.3. *Let V be an irreducible B -module with $J_B V \neq 0$. We define an A -module homomorphism*

$$\Phi : M \otimes_B V \rightarrow \text{Hom}_B(N, V)$$

by $\Phi(m \otimes v) : n \mapsto \varphi_B(n \otimes m)v$. Then, $\text{Ker}(\Phi)$ is the unique maximal proper A -submodule of $M \otimes_B V$.

In particular, $\text{Im}(\Phi) \simeq M \otimes_B V / \text{Ker}(\Phi)$ is an irreducible A -module, which we denote by $D_M^N V$. By the commutativity constraint for the Morita pair and $J_B V = V$, we have $J_A D_M^N V \neq 0$. Thus, we have a map $V \mapsto D_M^N V$ from the set of isomorphism classes of irreducible B -modules V with $J_B V \neq 0$ to the set of isomorphism classes of irreducible A -modules W with $J_A W \neq 0$. By interchanging the role of M and N , $W \mapsto D_N^M W$ gives a map in the opposite direction.

Next lemma by Mori implies that $D_M^N V = \text{Soc}(\text{Hom}_B(N, V))$. For the proof, he finds $b \in B$ and $n \in N$ which satisfy $\Phi(m \otimes v) = \varphi_A(mb \otimes n)f$ for any $m \otimes v \in M \otimes_B V$.

Lemma 5.4. *Let V and Φ be as above. For any nonzero element $f \in \text{Hom}_B(N, V)$, we have $\text{Im}(\Phi) \subseteq Af$.*

A main corollary of the above two lemmas is the following result by Mori. Mori informed us Müller's work [37], but Mori's argument is transparent and enough for our purposes.

Theorem 5.5. *Let (M, N) be a weak Morita pair. Then, $V \mapsto D_M^N V$ induces a bijective map between the set of isomorphism classes of irreducible B -modules V with $J_B V \neq 0$ and that of isomorphism classes of irreducible A -modules W with $J_A W \neq 0$.*

Proof. It suffices to show that $W \simeq D_M^N V$ if and only if $V \simeq D_N^M W$. Suppose that $W \simeq D_M^N V$. Then

$$0 \neq \text{Hom}_A(M \otimes_B V, W) \simeq \text{Hom}_B(V, \text{Hom}_A(M, W))$$

implies that V appears in $\text{Soc}(\text{Hom}_A(M, W)) \simeq D_N^M W$. If $V \simeq D_N^M W$, then

$$0 \neq \text{Hom}_B(N \otimes_A W, V) \simeq \text{Hom}_A(W, \text{Hom}_B(N, V))$$

implies that W appears in $\text{Soc}(\text{Hom}_B(N, V)) \simeq D_M^N V$. \square

Following those ideas by König-Xi and Mori, we shall define as follows.

Definition 5.6. Let (A, σ_A) be an \mathbf{k} -algebra with anti-involution. We call (A, σ_A) *generalized cellular* if there exists a sequence of σ_A -stable two-sided ideals

$$A = J_0 \supseteq J_1 \supseteq \cdots \supseteq J_r = 0$$

and a collection of \mathbf{k} -algebras with anti-involution (B_i, σ_{B_i}) and $(A/J_i, B_i)$ -bimodules C_i , for $1 \leq i \leq r$, such that (C_i, C_i^{op}) is a weak Morita pair with $J_{i-1}/J_i = \text{Im}(\varphi_{A/J_i})$, for $1 \leq i \leq r$.

Let us take all B_i to be \mathbf{k} and $\sigma_{\mathbf{k}}$ to be the identity map. Then, in each step, we have

$$\varphi_{A/J_i} : C_i \otimes_{\mathbf{k}} C_i^{\text{op}} \rightarrow A/J_i, \quad \text{and} \quad C_i^{\text{op}} \otimes_A C_i \rightarrow \mathbf{k}.$$

We denote the second map by $x \otimes y \mapsto \langle x, y \rangle$. As (C_i, C_i^{op}) is a weak Morita pair, we have

$$\varphi_{A/J_i}(x \otimes y)z = \langle y, z \rangle x, \quad \langle z, y \rangle x = \sigma_A(\varphi_{A/J_i}(y \otimes x))z.$$

If we further assume that φ_{A/J_i} commutes with anti-involutions, namely, if we assume

$$\varphi_{A/J_i}(y \otimes x) = \sigma_A(\varphi_{A/J_i}(x \otimes y)),$$

then

$$\langle z, y \rangle x = \sigma_A(\varphi_{A/J_i}(y \otimes x))z = \varphi_{A/J_i}(x \otimes y)z = \langle y, z \rangle x,$$

showing that the bilinear form is symmetric.¹ If we put one more assumption that φ_{A/J_i} are monomorphisms, then we have

$$C_i \otimes_R C_i^{\text{op}} \simeq J_{i-1}/J_i$$

¹If φ_{A/J_i} anti-commutes with anti-involutions, then the bilinear form is skew-symmetric.

and we reach the classical definition of cellular algebras. In fact, the commutativity constraint is known to hold in cellular algebras [36, Prop.2.9]. Hence, cellular algebras are generalized cellular algebras, and the reader can see that Theorem 5.5 is vast generalization of Graham-Lehrer's result on classification of irreducible modules.

5.2. Structure of $R^{\Lambda_0}(\beta)$. Any algebra has the trivial generalized cellular structure: we choose the cell bimodule C to be the algebra itself. Hence, our aim in this subsection is to propose a reasonable set of cell modules for $R^{\Lambda_0}(\beta)$, for $\beta \in \mathbf{Q}^+$ with $|\beta| = n$.

Set $A = R^{\Lambda_0}(\beta)$. Recall the dominance order on the set of shifted Young diagrams and the canonical tableaux T^λ from (1.1.1). We define idempotents $\{e_\lambda \mid \lambda \vdash n\}$ of A by

$$e_\lambda = e(\text{res}(T^\lambda)).$$

We fix a linear extension of the dominance order, e.g. the lexicographic order, and define $J^{<\lambda}, B^\lambda, C(\lambda)$, for $\lambda \vdash n$, as follows.

$$J^{<\lambda} = \sum_{\mu < \lambda} A e_\mu A, \quad B^\lambda = e_\lambda A e_\lambda / e_\lambda J^{<\lambda} e_\lambda, \quad \text{and} \quad C(\lambda) = A e_\lambda / J^{<\lambda} e_\lambda.$$

Then $C(\lambda)$ is an $(A/J^{<\lambda}, B^\lambda)$ -bimodule in the natural way.

Let σ_A be the anti-involution of A which is the identity on the set of the generators. Then, we have $C(\lambda)^{\text{op}} = e_\lambda A / e_\lambda J^{<\lambda}$. We define

$$\begin{aligned} \varphi_{A/J^{<\lambda}} &: C(\lambda) \otimes_{B^\lambda} C(\lambda)^{\text{op}} \longrightarrow A/J^{<\lambda}, \\ \varphi_{B^\lambda} &: C(\lambda)^{\text{op}} \otimes_A C(\lambda) \longrightarrow B^\lambda, \end{aligned}$$

by the maps induced by the multiplication on A . Then, we have

$$\text{Im}(\varphi_{A/J^{<\lambda}}) = A e_\lambda A + J^{<\lambda} / J^{<\lambda}.$$

We sort the two-sided ideals $J^{<\lambda}$ as

$$A = J_0 \supseteq J_1 \supseteq \cdots \supseteq J_r = 0.$$

We delete repetition whenever $J_i = J_{i+1}$ occurs.

Lemma 5.7. *The pair $(C(\lambda), C(\lambda)^{\text{op}})$ is a weak Morita pair, and $R^{\Lambda_0}(\beta)$ is generalized cellular with respect to the collection of those weak Morita pairs.*

Proof. The commutativity constraints obviously hold, because they are given by the product map in $R^{\Lambda_0}(\beta)$. \square

The generalized cellular structure we propose in the above looks natural, but we know little about the structure of cell modules. It might be an interesting question to determine the set of irreducible B^λ -modules V with $\text{Im}(\varphi_{B^\lambda})V \neq 0$.

REFERENCES

1. S. Ariki, Modular representations of Hecke algebras of classical type, *Sugaku Expositions* **20** (2007), 15–41. Translated from Japanese original: *Sugaku* **56** (2004), 113–136.
2. G. Benkart, S.-J. Kang, S.-j. Oh, E. Park, *Construction of Irreducible Representations over Khovanov-Lauda-Rouquier Algebras of Finite Classical Type*, arXiv:1108.1048.
3. J. Brundan and A. Kleshchev, *Hecke-Clifford superalgebras, crystals of type $A_{2\ell}^{(2)}$ and modular branching rules for \hat{S}_n* , Represent. Theory **5** (2001), 317–403.
4. ———, *Blocks of cyclotomic Hecke algebras and Khovanov-Lauda algebras*, Invent. Math. **178** (2009), 451–484,
5. J. Chuang and R. Rouquier, *Derived equivalences for symmetric groups and sl_2 -categorification*, Ann. of Math. (2) **167** (2008), no. 1, 245–298.
6. E. Date, M. Jimbo, M. Kashiwara, and T. Miwa, *Transformation groups for soliton equations. Euclidean Lie algebras and reduction of the KP hierarchy*, Publ. Res. Inst. Math. Sci. **18** (1982), no. 3, 1077–1110.
7. R. Dipper and G. D. James, *Blocks and idempotents of Hecke algebras of general linear groups*, Proc. London Math. Soc. (3), **54** (1987), 57–82.
8. K. Erdmann and D. K. Nakano, *Representation type of Hecke algebras of type A*, Trans. Amer. Math. Soc., **354** (2002), 275–285.
9. P. Gabriel and Ch. Riedtmann, *Group representations without groups*, Comment. Math. Helvetici **54** (1979) 240–287.
10. D. Hill, G. Melvin, and D. Mondragon, *Representations of quiver Hecke algebras via Lyndon bases*, J. Pure Appl. Algebra **216** (2012), no. 5, 1052–1079.
11. J. Hong and S.-J. Kang, *Introduction to Quantum Groups and Crystal Bases*, Graduate Studies in Mathematics, **42**. American Mathematical Society, Providence, RI, 2002.
12. G. D. James and A. Mathas, *A q -analogue of the Jantzen-Schaper theorem*, Proc. London Math. Soc. (3) (1997), **74**, 241–274.
13. G. J. Janusz, *Indecomposable Modules for finite groups*, Ann. of Math. **89** (1969), 209–241.
14. M. Jimbo and T. Miwa, *Solitons and infinite-dimensional Lie algebras*, Publ. Res. Inst. Math. Sci. **19** (1983), no. 3, 943–1001.
15. V. Kac, *Infinite-dimensional Lie algebras*, Third edition, Cambridge University Press, Cambridge, 1990.
16. S.-J. Kang, *Crystal bases for quantum affine algebras and combinatorics of Young walls*, Proc. London Math. Soc. (3) **86** (2003), no. 1, 29–69.
17. S.-J. Kang and M. Kashiwara, *Categorification of Highest Weight Modules via Khovanov-Lauda-Rouquier Algebras*, arXiv:1102.4677 (2011). to appear in Invent. Math.
18. S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima, and A. Nakayashiki, *Affine Crystals and Vertex Models*, Infinite Analysis, Part A,B(Kyoto, 1991), Adv. Ser. Math. Phys. **16**, World Scientific Publishing Co. Inc., River Edge, NJ, 1992, 449–484.
19. ———, *Perfect crystals of quantum affine Lie algebras*, Duke Math. J. **68** (1992), no. 3, 499–607.
20. S.-J. Kang and J.-H Kwon, *Fock space representations of quantum affine algebras and generalized Lascoux-Leclerc-Thibon algorithm*, J. Korean Math. Soc. **45** (2008), no. 4, 1135–1202.
21. S.-J. Kang and E. Park, *Irreducible modules over Khovanov-Lauda-Rouquier algebras of type A_n and semistandard tableaux*, J. Algebra, **339** (2011), no. 1, 223–251.

22. M. Kashiwara, *On crystal bases of the q -analogue of universal enveloping algebras*, Duke Math. J. **63** (1991), no. 2, 465–516.
23. ———, *The crystal base and Littelmann’s refined Demazure character formula*, Duke Math. J. **71** (1993), no. 3, 839–858.
24. ———, *Biadjointness in cyclic Khovanov-Lauda-Rouquier Algebras*, arXiv:1111.5898 (2011).
25. M. Kashiwara, T. Miwa, J.-U. H. Petersen, and C.M. Yung, *Perfect crystals and q -deformed Fock spaces*, Selecta Math. (N.S.) **2** (1996), no. 3, 415–499.
26. M. Khovanov and A. D. Lauda, *A diagrammatic approach to categorification of quantum groups I*, Represent. Theory **13** (2009), 309–347.
27. ———, *A diagrammatic approach to categorification of quantum groups II*, Trans. Amer. Math. Soc. **363** (2011), no. 5, 2685–2700.
28. A. Kleshchev and A. Ram, *Homogeneous Representations of Khovanov-Lauda Algebras*, J. Eur. Math. Soc. **12** (2010), 1293–1306.
29. ———, *Representations of Khovanov-Lauda-Rouquier Algebras and Combinatorics of Lyndon Words*, Math. Ann. **349** (2011), no. 4, 943–975.
30. S. König and C. C. Xi, *Affine cellular algebras*, Adv. Math. **229** (2012) 139–182.
31. H. Krause, *Stable equivalence preserves representation type*, Comment. Math. Helv., **72** (1997), 266–284.
32. A. D. Lauda, *An introduction to diagrammatic algebra and categorified quantum \mathfrak{sl}_2* , arXiv:1106.2128 (2011).
33. A. D. Lauda and M. Vazirani, *Crystals from categorified quantum groups*, Adv. Math. **228** (2011), no. 2, 803–861.
34. H. Lenzing, *Invariance of tameness under stable equivalence: Krause’s theorem*, in “Infinite Length Modules”, H. Krause and C. M. Ringel (eds.), Trends Math., (2000), 405–418.
35. S. Lyle and A. Mathas, *Blocks of cyclotomic Hecke algebras*, Adv. Math. **216** (2007), 854878.
36. A. Mathas, *Iwahori-Hecke Algebras and Schur Algebras of the Symmetric Group*, University Lecture Series, **15**. American Mathematical Society, Providence, RI, 1999.
37. B. Müller, *The quotient category of Morita context*, J. Algebra **28** (1974), 389–407.
38. W. K. Nicholson and J. F. Watters, *Morita context functors*, Math. Proc. Cambridge Philos. Soc. **103** (3) (1988), 399–408.
39. J. Rickard, *Derived categories and stable equivalence*, J. Pure and Appl. Alg. **61** (1989), 303–317.
40. C. M. Ringel, *The representation type of local algebras*, Representations of Algebras, Lecture Notes in Math. **488** (1975), 282–305.
41. R. Rouquier, *2 Kac-Moody algebras*, arXiv:0812.5023 (2008).
42. M. Varagnolo and E. Vasserot, *Canonical bases and KLR algebras*, J. Reine Angew. Math. **659** (2011), 67–100.

DEPARTMENT OF PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN

E-mail address: ariki@math.sci.osaka-u.ac.jp

DEPARTMENT OF PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN

E-mail address: pwy@cr.math.sci.osaka-u.ac.jp